

Extended AVF references (courtesy by Dr. G Beathard).

1. Dhingra, RK, Young, EW, Hulbert-Shearon, TE, et al. Type of vascular access and mortality in U.S. hemodialysis patients. *Kidney Int* 2001; 60:1443.
2. Polkinghorne, KR, McDonald, SP, Atkins, RC, Kerr, PG. Vascular access and all-cause mortality: a propensity score analysis. *J Am Soc Nephrol* 2004; 15:477.
3. Di Iorio, BR, Bellizzi, V, Cillo, N, et al. Vascular access for hemodialysis: the impact on morbidity and mortality. *J Nephrol* 2004; 17:19.
4. Ozeki, T, Shimizu, H, Fujita, Y, et al. The Type of Vascular Access and the Incidence of Mortality in Japanese Dialysis Patients. *Intern Med* 2017; 56:481.
5. Murad, MH, Elamin, MB, Sidawy, AN, et al. Autogenous versus prosthetic vascular access for hemodialysis: a systematic review and meta-analysis. *J Vasc Surg* 2008; 48:34S.
6. NKF-K/DOQI Clinical Practice Guidelines for Vascular Access: Update 2006. Guideline 8: Clinical outcome goals, 8.1 goals for access placement, 8.1.2 prevalent functional AVF placement rate.
7. Allon, M, Robbin, ML. Increasing arteriovenous fistulas in hemodialysis patients: problems and solutions. *Kidney Int* 2002; 62:1109.
8. Dember, LM, Beck, GJ, Allon, M, et al. Effect of clopidogrel on early failure of arteriovenous fistulas for hemodialysis: a randomized controlled trial. *JAMA* 2008; 299:2164.
9. Hodges, TC, Fillinger, MF, Zwolak, RM, et al. Longitudinal comparison of dialysis access methods: risk factors for failure. *J Vasc Surg* 1997; 26:1009.
10. Kalman, PG, Pope, M, Bhola, C, et al. A practical approach to vascular access for hemodialysis and predictors of success. *J Vasc Surg* 1999; 30:727.
11. Ascher, E, Gade, P, Hingorani, A, et al. Changes in the practice of Angio access surgery: impact of dialysis outcome and quality initiative recommendations. *J Vasc Surg* 2000; 31:84.
12. Hye, RJ, Wolf, YG. Ischemic monomelic neuropathy: an under-recognized complication of hemodialysis access. *Ann Vasc Surg* 1994; 8:578.
13. Lok, CE, Oliver, MJ. Overcoming barriers to arteriovenous fistula creation and use. *Semin Dial* 2003; 16:189.
14. Hernandez, T, Saudan, P, Berney, T, et al. Risk factors for early failure of native arteriovenous fistulas. *Nephron Clin Pract* 2005; 101:c39.
15. Patel, ST, Hughes, J, Mills, JL, Sr. Failure of arteriovenous fistula maturation: unintended

consequence of exceeding dialysis outcome quality Initiative guidelines for hemodialysis access. *J Vasc Surg* 2003; 38:439.

16. Lok, CE, Allon, M, Moist, L, et al. Risk equation determining unsuccessful cannulation events and failure to maturation in arteriovenous fistulas (REDUCE FTM I). *J Am Soc Nephrol* 2006; 17:3204.

17. Robbin, ML, Chamberlain, NE, Lockhart, ME, et al. Hemodialysis arteriovenous fistula maturity: US evaluation. *Radiology* 2002; 225:59.

18. Huijbregts, HJ, Bots, ML, Wittens, CH, et al. Hemodialysis arteriovenous fistula patency revisited: results of a prospective, multicenter initiative. *Clin J Am Soc Nephrol* 2008; 3:714.

19. Al-Jaishi, AA, Oliver, MJ, Thomas, SM, et al. Patency rates of the arteriovenous fistula for hemodialysis: a systematic review and meta-analysis. *Am J Kidney Dis* 2014; 63:464.

20. Lok, CE, Sontrop, JM, Tomlinson, G, et al. Cumulative patency of contemporary fistulas versus grafts (2000-2010). *Clin J Am Soc Nephrol* 2013; 8:810.

21. Woods, JD, Port, FK. The impact of vascular access for hemodialysis on patient morbidity and mortality. *Nephrol Dial Transplant* 1997; 12:657.

22. Dixon, BS, Novak, L, Fangman, J. Hemodialysis vascular access survival: upper-arm native

arteriovenous fistula. *Am J Kidney Dis* 2002; 39:92.

23. Perera, GB, Mueller, MP, Kubaska, SM, et al. Superiority of autogenous arteriovenous hemodialysis access: maintenance of function with fewer secondary interventions. *Ann Vasc Surg* 2004; 18:66.

24. Fitzgerald, JT, Schanzer, A, McVicar, JP, et al. Upper arm arteriovenous fistula versus forearm looped arteriovenous graft for hemodialysis access: a comparative analysis. *Ann Vasc Surg* 2005; 19:843.

25. Keuter, XH, De Smet, AA, Kessels, AG, et al. A randomized multicenter study of the outcome of brachial-basilic arteriovenous fistula and prosthetic brachial-antecubital forearm loop as vascular access for hemodialysis. *J Vasc Surg* 2008; 47:395.

26. Pisoni, RL, Young, EW, Dykstra, DM, et al. Vascular access use in Europe and the United States: results from the DOPPS. *Kidney Int* 2002; 61:305.

27. Huber, TS, Carter, JW, Carter, RL, Seeger, JM. Patency of autogenous and polytetrafluoroethylene upper extremity arteriovenous hemodialysis accesses: a systematic review. *J Vasc Surg* 2003; 38:1005.

28. Lee, T, Barker, J, Allon, M. Comparison of survival of upper arm arteriovenous fistulas and grafts after failed forearm fistula. *J Am Soc Nephrol* 2007; 18:1936.

29. Anel, RL, Yevzlin, AS, Ivanovich, P. Vascular access and patient outcomes in hemodialysis: questions answered in recent literature. *Artif Organs* 2003; 27:237.
30. Miles, AM. Vascular steal syndrome and ischemic monomelic neuropathy: two variants of upper limb ischemia after hemodialysis vascular access surgery. *Nephrol Dial Transplant* 1999; 14:297.
31. Miller, CD, Robbin, ML, Allon, M. Gender differences in outcomes of arteriovenous fistulas in hemodialysis patients. *Kidney Int* 2003; 63:346.
32. Tordoir, JH, Rooyens, P, Dammers, R, et al. Prospective evaluation of failure modes in autogenous radio cephalic wrist access for hemodialysis. *Nephrol Dial Transplant* 2003; 18:378.
33. Sidawy, AN, Gray, R, Besarab, A, et al. Recommended standards for reports dealing with arteriovenous hemodialysis accesses. *J Vasc Surg* 2002; 35:603.
34. Odland, MD, Kelly, PH, Ney, AL, et al. Management of dialysis-associated steal syndrome complicating upper extremity arteriovenous fistulas: use of intraoperative digital photoplethysmography. *Surgery* 1991; 110:664.
35. Lawson, JH, Glickman, MH, Ilzecki, M, et al. Bioengineered human acellular vessels for dialysis access in patients with end-stage renal disease: two phase 2 single-arm trials. *Lancet* 2016; 387:2026.
36. Gage, SM, Lawson, JH. Bioengineered hemodialysis access grafts. *J Vasc Access* 2017; 18:56.
37. Bonforte, G, Zerbi, S, Surian, M. The middle-arm fistula: A new native arteriovenous vascular access for hemodialysis patients. *Ann Vasc Surg* 2004; 18:448.
38. Wilbourn, AJ, Furlan, AJ, Hulley, W, Ruschhaupt, W. Ischemic monomelic neuropathy. *Neurology* 1983; 33:447.
39. Bonforte, G, Rossi, E, Auricchio, S, et al. The middle-arm fistula as a valuable surgical approach in patients with end-stage renal disease. *J Vasc Surg* 2010; 52:1551.
40. Jennings, WC, Mallios, A, Mushtaq, N. Proximal radial artery arteriovenous fistula for hemodialysis vascular access. *J Vasc Surg* 2018; 67:244.
41. Jennings, WC, Turman, MA, Taubman, KE. Arteriovenous fistulas for hemodialysis access in children and adolescents using the proximal radial artery inflow site. *J Pediatr Surg* 2009; 44:1377.
42. Jennings, WC, Landis, L, Taubman, KE, Parker, DE. Creating functional autogenous vascular access in older patients. *J Vasc Surg* 2011; 53:713.
43. Bolton, C, Dreidger, A, Lindsay, R. Ischemic neuropathy in uremic patients caused by bovine arteriovenous shunt. *J Neurol Neurosurg Psychiatry* 1974; 42:810.

44. Warren, DJ, Otieno, LS. Carpal tunnel syndrome in patients on intermittent haemodialysis. *Postgrad Med J* 1975; 51:450.
45. Holtmann, B, Anderson, CB. Carpal tunnel syndrome following vascular shunts for hemodialysis. *Arch Surg* 1977; 112:65.
46. Beathard, GA, Lok, CE, Glickman, MH, et al. Definitions and End Points for Interventional Studies for Arteriovenous Dialysis Access. *Clin J Am Soc Nephrol* 2017.
47. NKF-K/DOQI Clinical Practice Guidelines For Vascular Access: Update 2006. Guideline 1: Patient preparation for permanent hemodialysis access, 1.4 History and physical examination.
48. Beathard, G. *Complications of vascular access*, Marcel Dekker, Inc., New York, New York 2000.
49. Lok, CE, Foley, R. Vascular access morbidity and mortality: trends of the last decade. *Clin J Am Soc Nephrol* 2013; 8:1213.
50. Chang, VT, Hwang, SS, Feuerman, M, Kasimis, BS. Symptom and quality of life survey of medical oncology patients at a veteran's affairs medical center: a role for symptom assessment. *Cancer* 2000; 88:1175.
51. Weisbord, SD, Fried, LF, Arnold, RM, et al. Prevalence, severity, and importance of physical and emotional symptoms in chronic hemodialysis patients. *J Am Soc Nephrol* 2005; 16:2487.
52. Tonelli, M, Wiebe, N, Guthrie, B, et al. Comorbidity as a driver of adverse outcomes in people with chronic kidney disease. *Kidney Int* 2015; 88:859.
53. Crail, S, Walker, R, Brown, M, Renal Supportive Care working, g. Renal supportive and palliative care: position statement. *Nephrology (Carlton)* 2013; 18:393.
54. Cohen, LM, Germain, MJ, Woods, AL, et al. The family perspective of ESRD deaths. *Am J Kidney Dis* 2005; 45:154.
55. U.S. Renal Data System, *USRDS 2016 Annual Data Report*, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD.
56. Kalantar-Zadeh, K, Kovesdy, CP, Streja, E, et al. Transition of care from pre-dialysis prelude to renal replacement therapy: the blueprints of em
64. Burrowes, JD, Larive, B, Cockram, DB, et al. Effects of dietary intake, appetite, and eating habits on dialysis and non-dialysis treatment days in hemodialysis patients: cross-sectional results from the HEMO study. *J Ren Nutr* 2003; 13:191.
65. Carson, RC, Juszczak, M, Davenport, A, Burns, A. Is maximum conservative management an equivalent treatment option to dialysis for elderly patients with significant comorbid disease? *Clin J Am Soc Nephrol* 2009; 4:1611.

66. Lamping, DL, Constantinovici, N, Roderick, P, et al. Clinical outcomes, quality of life, and costs in the North Thames Dialysis Study of elderly people on dialysis: a prospective cohort study. *Lancet* 2000; 356:1543.
67. Foote, C, Kotwal, S, Gallagher, M, et al. Survival outcomes of supportive care versus dialysis therapies for elderly patients with end-stage kidney disease: A systematic review and meta-analysis. *Nephrology (Carlton)* 2016; 21:241.
68. Couchoud, C. Dialysis: Can we predict death in patients on dialysis? *Nat Rev Nephrol* 2010; 6:388.
69. Peterson, WJ, Barker, J, Allon, M. Disparities in fistula maturation persist despite preoperative vascular mapping. *Clin J Am Soc Nephrol* 2008; 3:437.
70. Lee, T, Ullah, A, Allon, M, et al. Decreased cumulative access survival in arteriovenous fistulas requiring interventions to promote maturation. *Clin J Am Soc Nephrol* 2011; 6:575.
71. Schild, AF, Prieto, J, Glenn, M, et al. Maturation and failure rates in a large series of arteriovenous dialysis access fistulas. *Vasc Endovascular Surg* 2004; 38:449.
72. Lok, CE, Oliver, MJ, Su, J, et al. Arteriovenous fistula outcomes in the era of the elderly dialysis population. *Kidney Int* 2005; 67:2462.
73. Chan, MR, Sanchez, RJ, Young, HN, Yevzlin, AS. Vascular access outcomes in the elderly hemodialysis population: A USRDS study. *Semin Dial* 2007; 20:606.
74. Beathard, GA, Spergel, LM. Hand ischemia associated with dialysis vascular access: an individualized access flow-based approach to therapy. *Semin Dial* 2013; 26:287.
75. Germain, MJ, Cohen, LM. Maintaining quality of life at the end of life in the end-stage renal disease population. *Adv Chronic Kidney Dis* 2008; 15:133.
76. O'Hare, AM, Allon, M, Kaufman, JS. Whether and when to refer patients for pre-dialysis AV fistula creation: complex decision making in the face of uncertainty. *Semin Dial* 2010; 23:452.
77. Keane, WF, Zhang, Z, Lyle, PA, et al. Risk scores for predicting outcomes in patients with type 2 diabetes and nephropathy: the RENAAL study. *Clin J Am Soc Nephrol* 2006; 1:761.
78. Keith, DS, Nichols, GA, Gullion, CM, et al. Longitudinal follow-up and outcomes among a population with chronic kidney disease in a large managed care organization. *Arch Intern Med* 2004; 164:659.
79. Lee, T, Barker, J, Allon, M. Tunneled catheters in hemodialysis patients: reasons and subsequent outcomes. *Am J Kidney Dis* 2005; 46:501.

80. Astor, BC, Eustace, JA, Powe, NR, et al. Timing of nephrologist referral and arteriovenous access use: the CHOICE Study. *Am J Kidney Dis* 2001; 38:494.
81. Lee, T, Barker, J, Allon, M. Associations with pre-dialysis vascular access management. *Am J Kidney Dis* 2004; 43:1008.
82. Vachharajani, TJ, Moossavi, S, Jordan, JR, et al. Re-evaluating the Fistula First Initiative in Octogenarians on Hemodialysis. *Clin J Am Soc Nephrol* 2011; 6:1663.
83. O'Hare, AM, Bertenthal, D, Walter, LC, et al. When to refer patients with chronic kidney disease for vascular access surgery: should age be a consideration? *Kidney Int* 2007; 71:555.
84. Richardson, AI, 2nd, Leake, A, Schmieder, GC, et al. Should fistulas really be first in the elderly patient? *J Vasc Access* 2009; 10:199.
85. NKF-K/DOQI Clinical Practice Guidelines For Vascular Access: Update 2006. Guideline 1: Patient preparation for permanent hemodialysis access.
86. Jindal, K, Chan, CT, Deziel, C, et al. Hemodialysis clinical practice guidelines for the Canadian Society of Nephrology. *J Am Soc Nephrol* 2006; 17: S1.
87. Tangri, N, Stevens, LA, Griffith, J, et al. A predictive model for progression of chronic kidney disease to kidney failure. *JAMA* 2011; 305:1553.
88. Tangri, N, Grams, ME, Levey, AS, et al. Multinational Assessment of Accuracy of Equations for Predicting Risk of Kidney Failure: A Meta-analysis. *JAMA* 2016; 315:164.
89. Cravedi, P, Remuzzi, G. Pathophysiology of proteinuria and its value as an outcome measure in chronic kidney disease. *Br J Clin Pharmacol* 2013; 76:516.
90. Levey, AS, Cattran, D, Friedman, A, et al. Proteinuria as a surrogate outcome in CKD: report of a scientific workshop sponsored by the National Kidney Foundation and the US Food and Drug Administration. *Am J Kidney Dis* 2009; 54:205.
91. Nitsch, D, Grams, M, Sang, Y, et al. Associations of estimated glomerular filtration rate and albuminuria with mortality and renal failure by sex: a meta-analysis. *BMJ* 2013; 346:f324.
92. Peeters, MJ, van Zuilen, AD, van den Brand, JA, et al. Validation of the kidney failure risk equation in European CKD patients. *Nephrol Dial Transplant* 2013; 28:1773.
93. Hingwala, J, Wojciechowski, P, Hiebert, B, et al. Risk-Based Triage for Nephrology Referrals Using the Kidney Failure Risk Equation. *Can J Kidney Health Dis* 2017; 4:2054358117722782.
94. Clase, CM, Kiberd, BA, Garg, AX. Relationship between glomerular filtration rate and the prevalence of metabolic abnormalities: results from the Third National Health and Nutrition Examination Survey (NHANES III). *Nephron Clin Pract* 2007; 105:c178.

95. Blantz, RC. Handing out grades for care in chronic kidney disease: nephrologists versus nonnephrologists. *Clin J Am Soc Nephrol* 2007; 2:193.
96. Kidney Disease Improving Global Outcomes (KDIGO) Work Group. KDIGO Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. Chapter 5: Referral to specialists and models of care. *Kidney International Supplements* 2013; 3:112.
97. Hiremath, S, Knoll, G, Weinstein, MC. Should the arteriovenous fistula be created before starting dialysis - a decision analytic approach. *PLoS One* 2011; 6: e28453.
98. Aitken, E, Thomson, P, Bainbridge, L, et al. A randomized controlled trial and cost-effectiveness analysis of early cannulation arteriovenous grafts versus tunneled central venous catheters in patients requiring urgent vascular access for hemodialysis. *J Vasc Surg* 2017; 65:766.
99. Ethier, J, Mendelssohn, DC, Elder, SJ, et al. Vascular access use and outcomes: an international perspective from the Dialysis Outcomes and Practice Patterns Study. *Nephrol Dial Transplant* 2008; 23:3219.
100. Pisoni, RL, Zepel, L, Port, FK, Robinson, BM. Trends in US Vascular Access Use, Patient Preferences, and Related Practices: An Update From the US DOPPS Practice Monitor With International Comparisons. *Am J Kidney Dis* 2015; 65:905.
101. Domenick Sridharan, N, Fish, L, Yu, L, et al. The associations of hemodialysis access type and access satisfaction with health-related quality of life. *J Vasc Surg* 2017.
102. Almasri, J, Alsawas, M, Mainou, M, et al. Outcomes of vascular access for hemodialysis: A systematic review and meta-analysis. *J Vasc Surg* 2016; 64:236.
103. DeSilva, R, Patibandla, B, Vin, Y, et al. Fistula first is not always the best strategy for the elderly. *J Am Soc Nephrol* 2013; 24.
104. Abbott, KC, Trespalacios, FC, Agodoa, LY. Arteriovenous fistula use and heart disease in long-term elderly hemodialysis patients: analysis of United States Renal Data System Dialysis Morbidity and Mortality Wave II. *J Nephrol* 2003; 16:822.
105. Malik, J, Tuka, V, Kasalova, Z, et al. Understanding the dialysis access steal syndrome. A review of the etiologies, diagnosis, prevention, and treatment strategies. *J Vasc Access* 2008; 9:155.
106. Spergel, LM, Ravani, P, Roy-Chaudhury, P, et al. Surgical salvage of the autogenous arteriovenous fistula (AVF). *J Nephrol* 2007; 20:388.
107. Jennings, WC, Brown, RE, Ruiz, C. Primary arteriovenous fistula inflow Proximalization for patients at high risk for dialysis access-associated ischemic steal syndrome. *J Vasc Surg* 2011; 54:554.
108. Hirai, M, Kawai, S. False positive and negative results in Allen test. *J Cardiovasc Surg (Torino)* 1980; 21:353.

109. Sidawy, AN, Spergel, LM, Besarab, A, et al. The Society for Vascular Surgery: clinical practice guidelines for the surgical placement and maintenance of arteriovenous hemodialysis access. *J Vasc Surg* 2008; 48:2S.
110. Tordoir, JH, Dammers, R, van der Sande, FM. Upper extremity ischemia and hemodialysis vascular access. *Eur J Vasc Endovasc Surg* 2004; 27:1.
111. van Hoek, F, Scheltinga, MR, Kouwenberg, I, et al. Stealing in hemodialysis patients depends on type of vascular access. *Eur J Vasc Endovasc Surg* 2006; 32:710.
112. Huijbregts, HJ, Bots, ML, Moll, FL, et al. Hospital specific aspects predominantly determine primary failure of hemodialysis arteriovenous fistulas. *J Vasc Surg* 2007; 45:962.
113. Farber, A, Imrey, PB, Huber, TS, et al. Multiple preoperative and intraoperative factors predict early fistula thrombosis in the Hemodialysis Fistula Maturation Study. *J Vasc Surg* 2016; 63:163.
114. Scheltinga, MR, van Hoek, F, Bruijninx, CM. Time of onset in hemodialysis access-induced distal ischemia (HAIDI) is related to the access type. *Nephrol Dial Transplant* 2009; 24:3198.
115. Georgakarakos, EI, Kapoulas, KC, Georgiadis, GS, et al. An overview of the hemodynamic aspects of the blood flow in the venous outflow tract of the arteriovenous fistula. *J Vasc Access* 2012; 13:271.
116. Sullivan, KL, Besarab, A, Dorrell, S, Moritz, MJ. The relationship between dialysis graft pressure and stenosis. *Invest Radiol* 1992; 27:352.
117. Saucy, F, Haesler, E, Haller, C, et al. Is intra-operative blood flow predictive for early failure of radiocephalic arteriovenous fistula? *Nephrol Dial Transplant* 2010; 25:862.
118. Kim, SY, Lee, JS, Kim, WO, et al. Evaluation of radial and ulnar blood flow after radial artery cannulation with 20- and 22-gauge cannula using duplex Doppler ultrasound. *Anaesthesia* 2012; 67:1138.
119. Asif, A, Roy-Chaudhury, P, Beathard, GA. Early arteriovenous fistula failure: a logical proposal for when and how to intervene. *Clin J Am Soc Nephrol* 2006; 1:332.
120. Girerd, X, London, G, Boutouyrie, P, et al. Remodeling of the radial artery in response to a chronic increase in shear stress. *Hypertension* 1996; 27:799.
121. van Gemert, MJ, Bruyninx, CM, Baggen, MJ. Shunt hemodynamics and extracorporeal dialysis: an electrical resistance network analysis. *Phys Med Biol* 1984; 29:219.
122. Ballermann, BJ, Dardik, A, Eng, E, Liu, A. Shear stress and the endothelium. *Kidney Int Suppl* 1998; 67:S100.

123. Joannides, R, Haefeli, WE, Linder, L, et al. Nitric oxide is responsible for flow-dependent dilatation of human peripheral conduit arteries in vivo. *Circulation* 1995; 91:1314.
124. Tohda, K, Masuda, H, Kawamura, K, Shozawa, T. Difference in dilatation between endothelium-preserved and -desquamated segments in the flow-loaded rat common carotid artery. *Arterioscler Thromb* 1992; 12:519.
125. Tronc, F, Wassef, M, Esposito, B, et al. Role of NO in flow-induced remodeling of the rabbit common carotid artery. *Arterioscler Thromb Vasc Biol* 1996; 16:1256.
126. Holtz, J, Forstermann, U, Pohl, U, et al. Flow-dependent, endothelium-mediated dilation of epicardial coronary arteries in conscious dogs: effects of cyclooxygenase inhibition. *J Cardiovasc Pharmacol* 1984; 6:1161.
127. Roy-Chaudhury, P, Spergel, LM, Besarab, A, et al. Biology of arteriovenous fistula failure. *J Nephrol* 2007; 20:150. Arteriovenous Fistula Beathard
128. Donnelly, SM, Marticorena, RM. When is a new fistula mature? The emerging science of fistula cannulation. *Semin Nephrol* 2012; 32:564.
129. Jaber, A, Muradali, D, Marticorena, RM, et al. Arteriovenous fistulas for hemodialysis: application of high-frequency US to assess vein wall morphology for cannulation readiness. *Radiology* 2011; 261:616.
130. Miller, VM, Burnett, JC, Jr. Modulation of NO and endothelin by chronic increases in blood flow in canine femoral arteries. *Am J Physiol* 1992; 263:H103.
131. Tronc, F, Mallat, Z, Lehoux, S, et al. Role of matrix metalloproteinases in blood flow-induced arterial enlargement: interaction with NO. *Arterioscler Thromb Vasc Biol* 2000; 20:E120.
132. Chan, CY, Chen, YS, Ma, MC, Chen, CF. Remodeling of experimental arteriovenous fistula with increased matrix metalloproteinase expression in rats. *J Vasc Surg* 2007; 45:804.
133. Chang, CJ, Chen, CC, Hsu, LA, et al. Degradation of the internal elastic laminae in vein grafts of rats with aortocaval fistulae: potential impact on graft vasculopathy. *Am J Pathol* 2009; 174:1837.
134. Dammers, R, Tordoir, JH, Kooman, JP, et al. The effect of flow changes on the arterial system proximal to an arteriovenous fistula for hemodialysis. *Ultrasound Med Biol* 2005; 31:1327.
135. Corpataux, JM, Haesler, E, Silacci, P, et al. Low-pressure environment and remodelling of the forearm vein in Brescia-Cimino haemodialysis access. *Nephrol Dial Transplant* 2002; 17:1057.
136. Dammers, R, Tordoir, JH, Welten, RJ, et al. The effect of chronic flow changes on brachial artery diameter and shear stress in arteriovenous fistulas for hemodialysis. *Int J Artif Organs* 2002; 25:124.

137. Lomonte, C, Casucci, F, Antonelli, M, et al. Is there a place for duplex screening of the brachial artery in the maturation of arteriovenous fistulas? *Semin Dial* 2005; 18:243.
138. Sigovan, M, Rayz, V, Gasper, W, et al. Vascular remodeling in autogenous arteriovenous fistulas by MRI and CFD. *Ann Biomed Eng* 2013; 41:657.
139. Beathard, GA, Arnold, P, Jackson, J, et al. Aggressive treatment of early fistula failure. *Kidney Int* 2003; 64:1487.
140. Zhu, YL, Ding, H, Fan, PL, et al. Predicting the maturity of haemodialysis arteriovenous fistulas with colour Doppler ultrasound: a single-centre study from China. *Clin Radiol* 2016; 71:576.
141. Wong, V, Ward, R, Taylor, J, et al. Factors associated with early failure of arteriovenous fistulae for HemoDialysis access. *Eur J Vasc Endovasc Surg* 1996; 12:207.
142. Lin, SL, Huang, CH, Chen, HS, et al. Effects of age and diabetes on blood flow rate and primary outcome of newly created hemodialysis arteriovenous fistulas. *Am J Nephrol* 1998; 18:96.
143. Malovrh, M. Native arteriovenous fistula: preoperative evaluation. *Am J Kidney Dis* 2002; 39:1218.
144. Ives, CL, Akoh, JA, George, J, et al. Pre-operative vessel mapping and early post-operative surveillance duplex scanning of arteriovenous fistulae. *J Vasc Access* 2009; 10:37.
145. Back, MR, Maynard, M, Winkler, A, Bandyk, DF. Expected flow parameters within hemodialysis access and selection for remedial intervention of nonmaturing conduits. *Vasc Endovascular Surg* 2008; 42:150.
146. Ferring, M, Henderson, J, Wilmlink, T. Accuracy of early postoperative clinical and ultrasound examination of arteriovenous fistulae to predict dialysis use. *J Vasc Access* 2014; 15:291.
147. Kamiya, A, Togawa, T. Adaptive regulation of wall shear stress to flow change in the canine carotid artery. *Am J Physiol* 1980; 239:H14.
148. Jemcov, TK. Morphologic and functional vessels characteristics assessed by ultrasonography for prediction of radio cephalic fistula maturation. *J Vasc Access* 2013; 14:356.
149. Besarab, A, Sherman, R. The relationship of recirculation to access blood flow. *Am J Kidney Dis* 1997; 29:223.
150. Wiese, P, Nonnast-Daniel, B. Colour Doppler ultrasound in dialysis access. *Nephrol Dial Transplant* 2004; 19:1956. Arteriovenous Fistula Beathard

151. Scaffaro, LA, Bettio, JA, Cavazzola, SA, et al. Maintenance of hemodialysis arteriovenous fistulas by an interventional strategy: clinical and duplex ultrasonographic surveillance followed by transluminal angioplasty. *J Ultrasound Med* 2009; 28:1159.
152. van Hooland, S, Malik, J. Hemodialysis vascular access ultrasonography: tips, tricks, pitfalls and a quiz. *J Vasc Access* 2010; 11:255.
153. Malovrh, M. Emerging role of vascular ultrasound in arteriovenous dialysis access, McGraw Hill Medica, New York 2012.
154. Chowdhury, S, Goss, D, Mistry, H, et al. Duplex ultrasound volumetric flow analysis before and after hemodialysis in patients with brachio-cephalic fistulae. *J Vasc Access* 2013; 14:342.
155. McCormack, LJ, Cauldwell, EW, Anson, BJ. Brachial and antebrachial arterial patterns; a study of 750 extremities. *Surg Gynecol Obstet* 1953; 96:43.
156. Rodriguez-Niedenfuhr, M, Vazquez, T, Nearn, L, et al. Variations of the arterial pattern in the upper limb revisited: a morphological and statistical study, with a review of the literature. *J Anat* 2001; 199:547.
157. Brescia, MJ, Cimino, JE, Appel, K, Hurwich, BJ. Chronic hemodialysis using venipuncture and a surgically created arteriovenous fistula. *N Engl J Med* 1966; 275:1089.
158. Mendelssohn, DC, Ethier, J, Elder, SJ, et al. Haemodialysis vascular access problems in Canada: results from the Dialysis Outcomes and Practice Patterns Study (DOPPS II). *Nephrol Dial Transplant* 2006; 21:721.
159. Woods, JD, Turenne, MN, Strawderman, RL, et al. Vascular access survival among incident hemodialysis patients in the United States. *Am J Kidney Dis* 1997; 30:50.
160. Chemla, ES, Morsy, MA. Is basilic vein transposition a real alternative to an arteriovenous bypass graft? A prospective study. *Semin Dial* 2008; 21:352.
161. Maya, ID, O'Neal, JC, Young, CJ, et al. Outcomes of brachiocephalic fistulas, transposed brachiobasilic fistulas, and upper arm grafts. *Clin J Am Soc Nephrol* 2009; 4:86.
162. Hakaim, AG, Nalbandian, M, Scott, T. Superior maturation and patency of primary brachiocephalic and transposed basilic vein arteriovenous fistulae in patients with diabetes. *J Vasc Surg* 1998; 27:154.
163. Moossavi, S, Tuttle, AB, Vachharajani, TJ, et al. Long-term outcomes of transposed basilic vein arteriovenous fistulae. *Hemodial Int* 2008; 12:80.
164. Silva, MB, Jr., Hobson, RW, 2nd, Pappas, PJ, et al. Vein transposition in the forearm for autogenous hemodialysis access. *J Vasc Surg* 1997; 26:981.

165. Silva, MB, Jr., Hobson, RW, 2nd, Pappas, PJ, et al. A strategy for increasing use of autogenous hemodialysis access procedures: impact of preoperative noninvasive evaluation. *J Vasc Surg* 1998; 27:302.
166. Ascher, E, Hingorani, A, Yorkovich, W. Techniques and out come after brachiocephalic and brachio basilic arteriovenous fistula creation, Lippincott, Williams & Wilkins, New York, New York 2002.
167. Moncef, G. Surgical revision of failing or thrombosed native arteriovenous fistulas: a single center experience. *Saudi J Kidney Dis Transpl* 2010; 21:258.
168. Turmel-Rodrigues, L, Mouton, A, Birmele, B, et al. Salvage of immature forearm fistulas for haemodialysis by interventional radiology. *Nephrol Dial Transplant* 2001; 16:2365.
169. Nassar, GM, Nguyen, B, Rhee, E, Achkar, K. Endovascular treatment of the "failing to mature" arteriovenous fistula. *Clin J Am Soc Nephrol* 2006; 1:275.
170. Poulain, F, Raynaud, A, Bourquelot, P, et al. Local thrombolysis and thrombo-aspiration in the treatment of acutely thrombosed arteriovenous hemodialysis fistulas. *Cardiovasc Intervent Radiol* 1991; 14:98.
171. Duijm, LE, Liem, YS, van der Rijt, RH, et al. Inflow stenoses in dysfunctional hemodialysis access fistulae and grafts. *Am J Kidney Dis* 2006; 48:98. Arteriovenous Fistula Beathard
172. Beathard, GA, Settle, SM, Shields, MW. Salvage of the nonfunctioning arteriovenous fistula. *Am J Kidney Dis* 1999; 33:910.
173. Manninen, HI, Kaukanen, E, Makinen, K, Karhapaa, P. Endovascular salvage of nonmaturing autogenous hemodialysis fistulas: comparison with endovascular therapy of failing mature fistulas. *J Vasc Interv Radiol* 2008; 19:870.
174. Napoli, M, Montinaro, A, Russo, F, et al. Early experiences of intraoperative ultrasound guided angioplasty of the arterial stenosis during upper limb arteriovenous fistula creation. *J Vasc Access* 2007; 8:97.
175. Falk, A. Maintenance and salvage of arteriovenous fistulas. *J Vasc Interv Radiol* 2006; 17:807.
176. Raynaud, A, Novelli, L, Bourquelot, P, et al. Low-flow maturation failure of distal accesses: Treatment by angioplasty of forearm arteries. *J Vasc Surg* 2009; 49:995.
177. Turmel-Rodrigues, L, Boutin, JM, Camiade, C, et al. Percutaneous dilation of the radial artery in non-maturing autogenous radial-cephalic fistulas for hemodialysis. *Nephrol Dial Transplant* 2009; 24:3782.
178. Pirozzi, N, Scrivano, J, Pirozzi, R, et al. Impaired maturation of arteriovenous fistula for hemodialysis due to forearm artery stenosis: percutaneous endovascular treatment. *J Vasc Access* 2017; 18:503.

179. Asif, A, Gadalean, FN, Merrill, D, et al. Inflow stenosis in arteriovenous fistulas and grafts: a multicenter prospective study. *Kidney Int* 2005; 67:1986.
180. Clark, TW, Cohen, RA, Kwak, A, et al. Salvage of nonmaturing native fistulas by using angioplasty. *Radiology* 2007; 242:286.
181. Shin, SW, Do, YS, Choo, SW, et al. Salvage of immature arteriovenous fistulas with percutaneous transluminal angioplasty. *Cardiovasc Intervent Radiol* 2005; 28:434.
182. Badero, OJ, Salifu, MO, Wasse, H, Work, J. Frequency of swing-segment stenosis in referred dialysis patients with angiographically documented lesions. *Am J Kidney Dis* 2008; 51:93.
183. Roy-Chaudhury, P, Lee, T, Woodle, B, et al. Balloon-assisted maturation (BAM) of the arteriovenous fistula: the good, the bad, and the ugly. *Semin Nephrol* 2012; 32:558.
184. Bourquelot, P, Raynaud, F, Pirozzi, N. Microsurgery in children for creation of arteriovenous fistulas in renal and non-renal diseases. *Ther Apher Dial* 2003; 7:498.
185. Hull, JE, Balakin, BV, Kellerman, BM, Wrolstad, DK. Computational fluid dynamic evaluation of the side-to-side anastomosis for arteriovenous fistula. *J Vasc Surg* 2013; 58:187.
186. Bharat, A, Jaenicke, M, Shenoy, S. A novel technique of vascular anastomosis to prevent juxta anastomotic stenosis following arteriovenous fistula creation. *J Vasc Surg* 2012; 55:274.
187. Sadaghianloo, N, Declémy, S, Jean-Baptiste, E, et al. Radial artery deviation and reimplantation inhibits venous juxta-anastomotic stenosis and increases primary patency of radial-cephalic fistulas for hemodialysis. *J Vasc Surg* 2016; 64:698.
188. Falk, A, Teodorescu, V, Lou, WY, et al. Treatment of "swing point stenoses" in hemodialysis arteriovenous fistulae. *Clin Nephrol* 2003; 60:35.
189. Turmel-Rodrigues, L, Pengloan, J, Baudin, S, et al. Treatment of stenosis and thrombosis in hemodialysis fistulas and grafts by interventional radiology. *Nephrol Dial Transplant* 2000; 15:2029.
190. Nassar, GM. Endovascular management of the "failing to mature" arteriovenous fistula. *Tech Vasc Interv Radiol* 2008; 11:175.
191. Castellan, L, Miotto, D, Savastano, S, et al. [The percutaneous transluminal angioplasty of Brescia-Cimino arteriovenous fistulae. An evaluation of the results]. *Radiol Med* 1994; 87:134.
192. Ortensia, A, Carosio, G, Iberti, M, et al. [Transluminal percutaneous angioplasty of stenosis of arteriovenous fistulas; 24-month follow-up]. *Minerva Urol Nefrol* 1998; 50:45.
193. Lay, JP, Ashleigh, RJ, Tranconi, L, et al. Result of angioplasty of brescia-cimino hemodialysis fistulae: medium-term follow-up. *Clin Radiol* 1998; 53:608.

194. Hingorani, A, Ascher, E, Kallakuri, S, et al. Impact of reintervention for failing upper-extremity arteriovenous autogenous access for hemodialysis. *J Vasc Surg* 2001; 34:1004.

195. Manninen, HI, Kaukanen, ET, Ikaheimo, R, et al. Brachial arterial access: endovascular treatment of failing Brescia-Cimino hemodialysis fistulas--initial success and long-term results. *Radiology* 2001; 218:711.

196. Clark, TW, Hirsch, DA, Jindal, KJ, et al. Outcome and prognostic factors of restenosis after

percutaneous treatment of native hemodialysis fistulas. *J Vasc Interv Radiol* 2002; 13:51.

197. Sugimoto, K, Higashino, T, Kuwata, Y, et al. Percutaneous transluminal angioplasty of malfunctioning Brescia-Cimino arteriovenous fistula: analysis of factors adversely affecting long-term patency. *Eur Radiol* 2003; 13:1615.

198. Rajan, DK, Bunston, S, Misra, S, et al. Dysfunctional autogenous hemodialysis fistulas: outcomes after angioplasty--are there clinical predictors of patency? *Radiology* 2004; 232:508.

199. Romero, A, Polo, JR, Garcia Morato, E, et al. Salvage of Angio access after late thrombosis of radio cephalic fistulas for hemodialysis. *Int Surg* 1986; 71:122.

200. Oakes, DD, Sherck, JP, Cobb, LF. Surgical salvage of failed radio cephalic arteriovenous fistulae: techniques and results in 29 patients. *Kidney Int* 1998; 53:480.

201. Murphy, GJ, White, SA, Nicholson, ML. Vascular access for hemodialysis. *Br J Surg* 2000; 87:1300.

202. Konner, K, Nonnast-Daniel, B, Ritz, E. The arteriovenous fistula. *J Am Soc Nephrol* 2003; 14:1669.

203. Napoli, M, Prudenzano, R, Russo, F, et al. Juxta-anastomotic stenosis of native arteriovenous fistulas: surgical treatment versus percutaneous transluminal angioplasty. *J Vasc Access* 2010; 11:346.

204. Long, B, Brichart, N, Lermusiaux, P, et al. Management of perianastomotic stenosis of direct wrist autogenous radial-cephalic arteriovenous accesses for dialysis. *J Vasc Surg* 2011; 53:108.

205. Tessitore, N, Mansueto, G, Lipari, G, et al. Endovascular versus surgical preemptive repair of forearm arteriovenous fistula juxta-anastomotic stenosis: analysis of data collected prospectively from 1999 to 2004. *Clin J Am Soc Nephrol* 2006; 1:448.

206. De Marco Garcia, LP, Davila-Santini, LR, Feng, Q, et al. Primary balloon angioplasty plus balloon angioplasty maturation to upgrade small-caliber veins (<3 mm) for arteriovenous fistulas. *J Vasc Surg* 2010; 52:139.

207. Veroux, P, Giaquinta, A, Tallarita, T, et al. Primary balloon angioplasty of small (≤ 2 mm) cephalic veins improves primary patency of arteriovenous fistulae and decreases reintervention rates. *J Vasc Surg* 2013; 57:131.
208. Miller, GA, Goel, N, Khariton, A, et al. Aggressive approach to salvage non-maturing arteriovenous fistulae: a retrospective study with follow-up. *J Vasc Access* 2009; 10:183.
209. DerDerian, T, Hingorani, A, Ascher, E, et al. To BAM or not to BAM? A closer look at balloon-assisted maturation. *Ann Vasc Surg* 2013; 27:104.
210. Chawla, A, DiRaimo, R, Panetta, TF. Balloon angioplasty to facilitate autogenous arteriovenous access maturation: a new paradigm for upgrading small-caliber veins, improved function, and surveillance. *Semin Vasc Surg* 2011; 24:82.
211. Faiyaz, R, Abreo, K, Zaman, F, et al. Salvage of poorly developed arteriovenous fistulae with percutaneous ligation of accessory veins. *Am J Kidney Dis* 2002; 39:824.
212. Planken, RN, Duijm, LE, Kessels, AG, et al. Accessory veins and radial-cephalic arteriovenous fistula non-maturation: a prospective analysis using contrast-enhanced magnetic resonance angiography. *J Vasc Access* 2007; 8:281.
213. Dixon, BS. Why don't fistulas mature? *Kidney Int* 2006; 70:1413.
214. McLafferty, RB. Techniques to enhance arteriovenous fistula maturation. *Perspect Vasc Surg Endovasc Ther* 2009; 21:41.
215. Nikam, M, Popuri, RK, Inaba, A, et al. Arteriovenous fistula failure: is there a role for accessory draining vein embolization? *J Vasc Access* 2012; 13:498.
216. Engstrom, BI, Grimm, LJ, Ronald, J, et al. Accessory veins in nonmaturing autogenous arteriovenous fistulae: analysis of anatomic features and impact on fistula maturation. *Semin Dial* 2015; 28:E30.
217. Haq, NU, Albaqumi, M. Accessory vein obliteration criteria for immature fistulae: a modest proposal for an old paradigm. *Semin Dial* 2014; 27:E51.
218. Usta, E, Elkrinawi, R, Salehi-Gilani, S, et al. Risk factors predicting the successful function and use of autogenous arteriovenous fistulae for hemodialysis. *Thorac Cardiovasc Surg* 2013; 61:438.
219. Ravani, P, Marcelli, D, Pecchini, P, Malberti, F. Early failure rates of arteriovenous fistulas for hemodialysis: evaluation of six-year activity. *J Vasc Access* 2001; 2:154.
220. Weber, CL, Djurdjev, O, Levin, A, Kiaii, M. Outcomes of vascular access creation prior to dialysis: building the case for early referral. *ASAIO J* 2009; 55:355.
221. Miller, GA, Hwang, W, Preddie, D, et al. Percutaneous salvage of thrombosed immature arteriovenous fistulas. *Semin Dial* 2011; 24:107.

222. Lazarides, MK, Staramos, DN, Kopadis, G, et al. Onset of arterial 'steal' following proximal Angio access: immediate and delayed types. *Nephrol Dial Transplant* 2003; 18:2387.
223. Pisoni, RL, Young, EW, Mapes, DL, et al. Vascular access use and outcomes in the U.S., Europe, and Japan: results from the Dialysis Outcomes and Practice Patterns Study. *Nephrol News Issues* 2003; 17:38.
224. Saran, R, Dykstra, DM, Pisoni, RL, et al. Timing of first cannulation and vascular access failure in hemodialysis: an analysis of practice patterns at dialysis facilities in the DOPPS. *Nephrol Dial Transplant* 2004; 19:2334.
225. Lee, T, Barker, J, Allon, M. Needle infiltration of arteriovenous fistulae in hemodialysis: risk factors and consequences. *Am J Kidney Dis* 2006; 47:1020.
226. Parisotto, MT, Schoder, VU, Miriunis, C, et al. Cannulation technique influences arteriovenous fistula and graft survival. *Kidney Int* 2014; 86:790.
227. Gallieni, M, Brenna, I, Brunini, F, et al. Which cannulation technique for which patient. *J Vasc Access* 2014; 15 Suppl 7:S85.
228. Twardowski, Z, Lebek, R, Kubara, H. [6-year experience with the creation and use of internal arteriovenous fistulae in patients treated with repeated hemodialysis]. *Pol Arch Med Wewn* 1977; 57:205.
229. Ward, J, Shaw, K, Davenport, A. Patients' perspectives of constant-site (buttonhole) cannulation for hemodialysis access. *Nephron Clin Pract* 2010; 116:c123.
230. Ball, LK. The buttonhole technique for arteriovenous fistula cannulation. *Nephrol Nurs J* 2006; 33:299.
231. Twardowski, Z. Constant site (buttonhole) method of needle insertion for hemodialysis. *Dial Transplant* 1995; 24:559.
232. van Loon, MM, Goovaerts, T, Kessels, AG, et al. Buttonhole needling of haemodialysis arteriovenous fistulae results in less complications and interventions compared to the rope-ladder technique. *Nephrol Dial Transplant* 2010; 25:225.
233. MacRae, JM, Ahmed, SB, Atkar, R, Hemmelgarn, BR. A randomized trial comparing buttonholes with rope ladder needling in conventional hemodialysis patients. *Clin J Am Soc Nephrol* 2012; 7:1632.
234. Nesrallah, GE, Cuerden, M, Wong, JH, Pierratos, A. Staphylococcus aureus bacteremia and buttonhole cannulation: long-term safety and efficacy of mupirocin prophylaxis. *Clin J Am Soc Nephrol* 2010; 5:1047.
235. Van Eps, CL, Jones, M, Ng, T, et al. The impact of extended-hours home hemodialysis and buttonhole cannulation technique on hospitalization rates for septic events related to dialysis access. *Hemodial Int* 2010; 14:451.

236. Verhallen, AM, Kooistra, MP, van Jaarsveld, BC. Cannulating hemodialysis: rope-ladder or buttonhole technique? *Nephrol Dial Transplant* 2007; 22:2601.
237. Doss, S, Schiller, B, Moran, J. Buttonhole cannulation--an unexpected outcome. *Nephrol Nurs J* 2008; 35:417.
238. Ludlow, V. Buttonhole cannulation in hemodialysis: improved outcomes and increased expense--is it worth it? *CANNT J* 2010; 20:29.
239. Murcutt, G. Buttonhole cannulation: should this become the default technique for dialysis patients with native fistulas? Summary of the EDTNA/ERCA Journal Club discussion Autumn 2007. *J Ren Care* 2008;34:101.
240. Labriola, L, Crott, R, Desmet, C, et al. Infectious complications following conversion to buttonhole cannulation of native arteriovenous fistulas: a quality improvement report. *Am J Kidney Dis* 2011; 57:442.
241. Labriola, L, Jadoul, M. Infectious complications following conversion to buttonhole cannulation. *Clin Nephrol* 2011; 76:423; author reply 424.
242. Muir, CA, Kotwal, SS, Hawley, CM, et al. Buttonhole cannulation and clinical outcomes in a home hemodialysis cohort and systematic review. *Clin J Am Soc Nephrol* 2014; 9:110.
243. Silva Gdos, S, Silva, RA, Nicolino, AM, et al. Initial experience with the buttonhole technique in a Brazilian hemodialysis center. *J Bras Nefrol* 2010; 32:257.
244. Fistula First. Change concept 4.
http://www.fistulafirst.org/Professionals/FFBIChange_Concepts.aspx.
245. Sivanesan, S, How, TV, Bakran, A. Sites of stenosis in AV fistulae for haemodialysis access. *Nephrol Dial Transplant* 1999; 14:118.
246. Hammes, M, Funaki, B, Coe, FL. Cephalic arch stenosis in patients with fistula access for hemodialysis: relationship to diabetes and thrombosis. *Hemodial Int* 2008; 12:85.
247. Rajan, DK, Clark, TW, Patel, NK, et al. Prevalence and treatment of cephalic arch stenosis in dysfunctional autogenous hemodialysis fistulas. *J Vasc Interv Radiol* 2003; 14:567.
248. Beaulieu, MC, Gabana, C, Rose, C, et al. Stenosis at the area of transposition - an under-recognized complication of transposed brachio basilic fistulas. *J Vasc Access* 2007; 8:268.
249. Oliver, MJ, McCann, RL, Indridason, OS, et al. Comparison of transposed brachio basilic fistulas to upper arm grafts and brachiocephalic fistulas. *Kidney Int* 2001; 60:1532.

250. Chang, CJ, Ko, PJ, Hsu, LA, et al. Highly increased cell proliferation activity in the restenosis hemodialysis vascular access after percutaneous transluminal angioplasty: implication in prevention of restenosis. *Am J Kidney Dis* 2004; 43:74.
251. Besarab, A, Asif, A, Roy-Chaudhury, P, et al. The native arteriovenous fistula in 2007. Surveillance and monitoring. *J Nephrol* 2007; 20:656.
252. Paulson, WD, Moist, L, Lok, CE. Vascular access surveillance: an ongoing controversy. *Kidney Int* 2012; 81:132.
253. Heerwagen, ST, Hansen, MA, Schroeder, TV, et al. Endovascular treatment of hemodialysis arteriovenous fistulas: is immediate post-interventional blood flow a predictor of patency. *J Vasc Access* 2012; 13:315.
254. Neuen, BL, Gunnarsson, R, Webster, AC, et al. Predictors of patency after balloon angioplasty in hemodialysis fistulas: a systematic review. *J Vasc Interv Radiol* 2014; 25:917.
255. Fahrtash, F, Kairaitis, L, Gruenewald, S, et al. Defining a significant stenosis in an autologous radiocephalic arteriovenous fistula for hemodialysis. *Semin Dial* 2011; 24:231.
256. Fluck, R, Kumwenda, M, hemodialysis, Racpgovaf. Renal association clinical practice guidelines on vascular access for hemodialysis *Nephron Clin Pract* 2011:118:c225.
257. Tessitore, N, Bedogna, V, Poli, A, et al. Should current criteria for detecting and repairing arteriovenous fistula stenosis be reconsidered? Interim analysis of a randomized controlled trial. *Nephrol Dial Transplant* 2014; 29:179.
258. Beathard, G. Complications of vascular access. In: *Complications of Dialysis – Recognition and Management.*, Lameire, N, Mehta, R (Eds), Marcel Dekker, Inc, New York 2000. p. 1.
259. Beathard, GA. Percutaneous transvenous angioplasty in the treatment of vascular access stenosis. *Kidney Int* 1992; 42:1390.
260. Mori, Y, Horikawa, K, Sato, K, et al. Stenotic lesions in vascular access: treatment with transluminal angioplasty using high-pressure balloons. *Intern Med* 1994; 33:284.
261. Beathard, GA. The treatment of vascular access graft dysfunction: a nephrologist's view and experience. *Adv Ren Replace Ther* 1994; 1:131.
262. Kanterman, RY, Vesely, TM, Pilgram, TK, et al. Dialysis access grafts: anatomic location of venous stenosis and results of angioplasty. *Radiology* 1995; 195:135.
263. Lumsden, AB, MacDonald, MJ, Kikeri, DK, et al. Hemodialysis access graft stenosis: percutaneous transluminal angioplasty. *J Surg Res* 1997; 68:181.

264. Longwitz, D, Pham, TH, Heckemann, RG, Hecking, E. [Angioplasty in the stenosed hemodialysis shunt: experiences with 100 patients and 166 interventions]. *Rofo* 1998; 169:68.
265. Lilly, RZ, Carlton, D, Barker, J, et al. Predictors of arteriovenous graft patency after radiologic intervention in hemodialysis patients. *Am J Kidney Dis* 2001; 37:945.
266. Beathard, GA, Litchfield, T, Physician Operators Forum of Rms Lifeline, I. Effectiveness, and safety of dialysis vascular access procedures performed by interventional nephrologists. *Kidney Int* 2004; 66:1622.
267. Suzuki, K, Narimatsu, Y, Ido, K, et al. [Percutaneous transluminal angioplasty for stenotic dialysis arterio-venous fistulas]. *Nihon Igaku Hoshasen Gakkai Zasshi* 1992; 52:344.
268. Maeda, K, Furukawa, A, Yamasaki, M, Murata, K. Percutaneous transluminal angioplasty for Brescia-Cimino hemodialysis fistula dysfunction: technical success rate, patency rate and factors that influence the results. *Eur J Radiol* 2005; 54:426.
269. Wu, CC, Lin, MC, Pu, SY, et al. Comparison of cutting balloon versus high-pressure balloon angioplasty for resistant venous stenoses of native hemodialysis fistulas. *J Vasc Interv Radiol* 2008; 19:877.
270. Yang, TY, Cheng, HW, Weng, HH, et al. Percutaneous transluminal angioplasty for radial-cephalic fistulae with stenosis at the arteriovenous junction. *Am J Med Sci* 2012; 343:435.
271. Glanz, S, Gordon, DH, Lipkowitz, GS, et al. Axillary and subclavian vein stenosis: percutaneous angioplasty. *Radiology* 1988; 168:371.
272. Kovalik, EC, Newman, GE, Suhocki, P, et al. Correction of central venous stenoses: use of angioplasty and vascular Wallstents. *Kidney Int* 1994; 45:1177.
273. Quinn, SF, Schuman, ES, Demlow, TA, et al. Percutaneous transluminal angioplasty versus endovascular stent placement in the treatment of venous stenoses in patients undergoing hemodialysis: intermediate results. *J Vasc Interv Radiol* 1995; 6:851.
274. Dammers, R, de Haan, MW, Planken, NR, et al. Central vein obstruction in hemodialysis patients: results of radiological and surgical intervention. *Eur J Vasc Endovasc Surg* 2003; 26:317.
275. Surowiec, SM, Fegley, AJ, Tanski, WJ, et al. Endovascular management of central venous stenoses in the hemodialysis patient: results of percutaneous therapy. *Vasc Endovascular Surg* 2004; 38:349.
276. Bakken, AM, Protack, CD, Saad, WE, et al. Long-term outcomes of primary angioplasty and primary stenting of central venous stenosis in hemodialysis patients. *J Vasc Surg* 2007; 45:776.

277. Haage, P, Vorwerk, D, Wildberger, JE, et al. Percutaneous treatment of thrombosed primary arteriovenous hemodialysis access fistulae. *Kidney Int* 2000; 57:1169.
278. Chang, CJ, Ko, YS, Ko, PJ, et al. Thrombosed arteriovenous fistula for hemodialysis access is characterized by a marked inflammatory activity. *Kidney Int* 2005; 68:1312.
279. Jain, G, Maya, ID, Allon, M. Outcomes of percutaneous mechanical thrombectomy of arteriovenous fistulas in hemodialysis patients. *Semin Dial* 2008; 21:581.
280. Schon, D, Mishler, R. Salvage of occluded autologous arteriovenous fistulae. *Am J Kidney Dis* 2000; 36:804.
281. Schon, D, Mishler, R. Pharmacochemical thrombolysis of natural vein fistulas: reduced dose of TPA and long-term follow-up. *Semin Dial* 2003; 16:272.
282. Turmel-Rodrigues, L, Pengloan, J, Rodrigue, H, et al. Treatment of failed native arteriovenous fistulae for hemodialysis by interventional radiology. *Kidney Int* 2000; 57:1124.
283. Huang, HL, Chen, CC, Chang, SH, et al. Combination of duplex ultrasound-guided manual declotting and percutaneous transluminal angioplasty in thrombosed native dialysis fistulas. *Ren Fail* 2005; 27:713.
284. Shatsky, JB, Berns, JS, Clark, TW, et al. Single-center experience with the Arrow-Treterotola Percutaneous Thrombectomy Device in the management of thrombosed native dialysis fistulas. *J Vasc Interv Radiol* 2005; 16:1605.
285. Padberg, FT, Jr., Calligaro, KD, Sidawy, AN. Complications of arteriovenous hemodialysis access: recognition and management. *J Vasc Surg* 2008; 48:55S.
286. Mennes, PA, Gilula, LA, Anderson, CB, et al. Complications associated with arteriovenous fistulas in patients undergoing chronic hemodialysis. *Arch Intern Med* 1978; 138:1117.
287. Derakhshanfar, A, Gholyaf, M, Niayesh, A, Bahiraii, S. Assessment of frequency of complications of arterio venous fistula in patients on dialysis: a two-year single center study from Iran. *Saudi J Kidney Dis Transpl* 2009; 20:872.
288. Pasklinsky, G, Meisner, RJ, Labropoulos, N, et al. Management of true aneurysms of hemodialysis access fistulas. *J Vasc Surg* 2011; 53:1291.
289. Salahi, H, Fazelzadeh, A, Mehdizadeh, A, et al. Complications of arteriovenous fistula in dialysis patients. *Transplant Proc* 2006; 38:1261.
290. Balaz, P, Bjorck, M. True aneurysm in autologous hemodialysis fistulae: definitions, classification and indications for treatment. *J Vasc Access* 2015; 16:446.
291. Inston, N, Mistry, H, Gilbert, J, et al. Aneurysms in vascular access: state of the art and future developments. *J Vasc Access* 2017; 18:464.

292. Valenti, D, Mistry, H, Stephenson, M. A novel classification system for autogenous arteriovenous fistula aneurysms in renal access patients. *Vasc Endovascular Surg* 2014; 48:491.
293. Unnikrishnan, S, Huynh, TN, Brott, BC, et al. Turbulent flow evaluation of the venous needle during hemodialysis. *J Biomech Eng* 2005; 127:1141.
294. Huynh, TN, Chacko, BK, Teng, X, et al. Effects of venous needle turbulence during ex vivo hemodialysis on endothelial morphology and nitric oxide formation. *J Biomech* 2007; 40:2158.
295. Rajput, A, Rajan, DK, Simons, ME, et al. Venous aneurysms in autogenous hemodialysis fistulas: is there an association with venous outflow stenosis. *J Vasc Access* 2013; 14:126.
296. Georgakarakos, EI, Xenakis, A, Kapoulas, KC, et al. A computational study of the factors influencing pressure in arteriovenous fistulae venous aneurysms. *J Vasc Access* 2013; 14:325.
297. Cunnane, CV, Cunnane, EM, Walsh, MT. A Review of the Hemodynamic Factors Believed to Contribute to Vascular Access Dysfunction. *Cardiovasc Eng Technol* 2017; 8:280.
298. Georgiadis, GS, Nikolopoulos, E, Papanas, N, et al. A hybrid approach to salvage a failing longstanding autogenous aneurysmal fistula in a hemodialysis patient. *Int J Artif Organs* 2010; 33:819.
299. Pierce, GE, Thomas, JH, Fenton, JR. Novel repair of venous aneurysms secondary to arteriovenous dialysis fistulae. *Vasc Endovascular Surg* 2007; 41:55.
300. Georgiadis, GS, Lazarides, MK, Panagoutsos, SA, et al. Surgical revision of complicated false and true vascular access-related aneurysms. *J Vasc Surg* 2008; 47:1284.
301. Berard, X, Brizzi, V, Mayeux, S, et al. Salvage treatment for venous aneurysm complicating vascular access arteriovenous fistula: use of an Arteriovenous Fistula
Beathard
302. Georgiadis, GS, Lazarides, MK, Lambidis, CD, et al. Use of short PTFE segments (<6 cm) compares favorably with pure autologous repair in failing or thrombosed native arteriovenous fistulas. *J Vasc Surg* 2005; 41:76.
303. Wytrzes, L, Markley, HG, Fisher, M, Alfred, HJ. Brachial neuropathy after brachial artery-antecubital vein shunts for chronic hemodialysis. *Neurology* 1987; 37:1398.
304. Redfern, AB, Zimmerman, NB. Neurologic and ischemic complications of upper extremity vascular access for dialysis. *J Hand Surg Am* 1995; 20:199.

305. Isoda, S, Kajiwara, H, Kondo, J, Matsumoto, A. Banding a hemodialysis arteriovenous fistula to decrease blood flow and resolve high output cardiac failure: report of a case. *Surg Today* 1994; 24:734.
306. Young, PR, Jr., Rohr, MS, Marterre, WF, Jr. High-output cardiac failure secondary to a brachiocephalic arteriovenous hemodialysis fistula: two cases. *Am Surg* 1998; 64:239.
307. MacRae, JM, Pandeya, S, Humen, DP, et al. Arteriovenous fistula-associated high-output cardiac failure: a review of mechanisms. *Am J Kidney Dis* 2004; 43:e17.
308. Wasse, H, Singapuri, MS. High-output heart failure: how to define it, when to treat it, and how to treat it. *Semin Nephrol* 2012; 32:551.
309. Basile, C, Lomonte, C, Vernaglione, L, et al. The relationship between the flow of arteriovenous fistula and cardiac output in hemodialysis patients. *Nephrol Dial Transplant* 2008; 23:282.
310. Begin, V, Ethier, J, Dumont, M, Leblanc, M. Prospective evaluation of the intra-access flow of recently created native arteriovenous fistulae. *Am J Kidney Dis* 2002; 40:1277.
311. Zanow, J, Petzold, K, Petzold, M, et al. Flow reduction in high-flow arteriovenous access using intraoperative flow monitoring. *J Vasc Surg* 2006; 44:1273.
312. Wijnen, E, Keuter, XH, Planken, NR, et al. The relation between vascular access flow and different types of vascular access with systemic hemodynamics in hemodialysis patients. *Artif Organs* 2005; 29:960.
313. Aitken, E, Kerr, D, Geddes, C, et al. Cardiovascular changes occurring with occlusion of a mature arteriovenous fistula. *J Vasc Access* 2015; 16:459.
314. Coffey, JP, Woywodt, A, Hill, JC. SPECT MIBI imaging for cardiac output and index in end stage renal disease. *Hemodial Int* 2011; 15:320.
315. Engelberts, I, Tordoir, JH, Boon, ES, Schreij, G. High-output cardiac failure due to excessive shunting in a hemodialysis access fistula: an easily overlooked diagnosis. *Am J Nephrol* 1995; 15:323.
316. Stern, AB, Klemmer, PJ. High-output heart failure is secondary to arteriovenous fistula. *Hemodial Int* 2011.
317. Borlaug, B, Beathard, G. Evaluation and management of heart failure caused by hemodialysis arteriovenous access. In: *UpToDate* 2018.
318. Reddy, YN, Melenovsky, V, Redfield, MM, et al. High-Output Heart Failure: A 15-Year Experience. *J Am Coll Cardiol* 2016; 68:473.
319. Anand, IS, Florea, VG. High Output Cardiac Failure. *Curr Treat Options Cardiovasc Med* 2001; 3:151.

320. Wohlfahrt, P, Rokosny, S, Melenovsky, V, et al. Cardiac remodeling after reduction of high-flow arteriovenous fistulas in end-stage renal disease. *Hypertens Res* 2016; 39:654.
321. Gkotsis, G, Jennings, WC, Malik, J, et al. Treatment of High Flow Arteriovenous Fistulas after Successful Renal Transplant Using a Simple Precision Banding Technique. *Ann Vasc Surg* 2016; 31:85.
322. Velez-Roa, S, Neubauer, J, Wissing, M, et al. Acute arterio-venous fistula occlusion decreases sympathetic activity and improves baroreflex control in kidney transplanted patients. *Nephrol Dial Transplant* 2004; 19:1606.
323. Bos, WJ, Zietse, R, Wesseling, KH, Westerhof, N. Effects of arteriovenous fistulas on cardiac oxygen supply and demand. *Kidney Int* 1999; 55:2049.
324. Unger, P, Wissing, KM, de Pauw, L, et al. Reduction of left ventricular diameter and mass after surgical arteriovenous fistula closure in renal transplant recipients. *Transplantation* 2002; 74:73.
325. van Hoek, F, Scheltinga, M, Luirink, M, et al. Banding of hemodialysis access to treat hand ischemia or cardiac overload. *Semin Dial* 2009; 22:204.
326. Miller, GA, Goel, N, Friedman, A, et al. The MILLER banding procedure is an effective method for treating dialysis-associated steal syndrome. *Kidney Int* 2010; 77:359.
327. Chemla, ES, Morsy, M, Anderson, L, Whitmore, A. Inflow reduction by distalization of anastomosis treats efficiently high-inflow high-cardiac output vascular access for hemodialysis. *Semin Dial* 2007; 20:68.
328. Parmar, CD, Chieng, G, Abraham, KA, et al. Revision using distal inflow for treatment of heart failure secondary to arteriovenous fistula for hemodialysis. *J Vasc Access* 2009; 10:62.
329. Bourquelot, P. Access flow reduction for cardiac failure. *J Vasc Access* 2016; 17 Suppl 1:S60.
330. Taylor, B, Sigley, RD, May, KJ. Fate of infected and eroded hemodialysis grafts and autogenous fistulas. *Am J Surg* 1993; 165:632.
331. Fong, IW, Capellan, JM, Simbul, M, Angel, J. Infection of arterio-venous fistulas created for chronic hemodialysis. *Scand J Infect Dis* 1993; 25:215.
332. Churchill, DN, Taylor, DW, Cook, RJ, et al. Canadian Hemodialysis Morbidity Study. *Am J Kidney Dis* 1992; 19:214.
333. Ravani, P, Palmer, SC, Oliver, MJ, et al. Associations between hemodialysis access type and clinical outcomes: a systematic review. *J Am Soc Nephrol* 2013; 24:465.

334. Kornfield, ZN, Kwak, A, Soulen, MC, et al. Incidence and management of percutaneous transluminal angioplasty-induced venous rupture in the "fistula first" era. *J Vasc Interv Radiol* 2009; 20:744.
335. Pappas, JN, Vesely, TM. Vascular rupture during angioplasty of hemodialysis raft-related stenoses. *J Vasc Access* 2002; 3:120.
336. Beathard, GA, Urbanes, A, Litchfield, T. Changes in the Profile of Endovascular Procedures Performed in Freestanding Dialysis Access Centers over 15 Years. *Clin J Am Soc Nephrol* 2017; 12:779.
337. Beathard, GA, Urbanes, A, Litchfield, T. The classification of procedure-related complications--a fresh approach. *Semin Dial* 2006; 19:527.
338. Beathard, GA. Strategy for maximizing the use of arteriovenous fistulae. *Semin Dial* 2000; 13:291.
339. Beathard, GA. Interventionalist's role in identifying candidates for secondary fistulas. *Semin Dial* 2004; 17:233.
340. Nguyen, VD, Treat, L, Griffith, C, Robinson, K. Creation of secondary AV fistulas from failed hemodialysis grafts: the role of routine vein mapping. *J Vasc Access* 2007; 8:91.
341. Work, J. How can the use of arteriovenous fistulas be increased? *Semin Dial* 2003; 16:220.
342. Asif, A, Unger, SW, Briones, P, et al. Creation of secondary arteriovenous fistulas: maximizing fistulas in prevalent hemodialysis patients. *Semin Dial* 2005; 18:420.
343. Jennings, WC. Creating arteriovenous fistulas in 132 consecutive patients: exploiting the proximal radial artery arteriovenous fistula: reliable, safe, and simple forearm and upper arm hemodialysis access. *Arch Surg* 2006; 141:27.
344. Robbin, ML, Gallichio, MH, Deierhoi, MH, et al. US vascular mapping before hemodialysis access placement. *Radiology* 2000; 217:83.
345. Allon, M, Lockhart, ME, Lilly, RZ, et al. Effect of preoperative sonographic mapping on vascular access outcomes in hemodialysis patients. *Kidney Int* 2001; 60:2013.
346. Ilhan, G, Esi, E, Bozok, S, et al. The clinical utility of vascular mapping with Doppler ultrasound prior to arteriovenous fistula construction for hemodialysis access. *J Vasc Access* 2013; 14:83.
347. Smith, GE, Samuel, N, Khan, J, et al. Targeted duplex ultrasound in a one-stop dialysis vascular access assessment clinic. *Ann Vasc Surg* 2011; 25:1099.
348. Anaya-Ayala, JE, Younes, HK, Kaiser, CL, et al. Prevalence of variant brachial-basilic vein anatomy and implications for vascular access planning. *J Vasc Surg* 2011; 53:720.

349. Kaiser, CL, Anaya-Ayala, JE, Ismail, N, et al. Unrecognized basilic vein variation leading to complication during basilic vein transposition arteriovenous fistula creation: case report and implications for access planning. *Eur J Vasc Endovasc Surg* 2010; 39:627.
350. Kurella, M, Covinsky, KE, Collins, AJ, Chertow, GM. Octogenarians and nonagenarians starting dialysis in the United States. *Ann Intern Med* 2007; 146:177.
351. Lee, T, Thamer, M, Zhang, Y, et al. Outcomes of Elderly Patients after Pre-dialysis Vascular Access Creation. *J Am Soc Nephrol* 2015; 26:3133.
352. Lazarides, MK, Georgiadis, GS, Antoniou, GA, Staramos, DN. A meta-analysis of dialysis access outcome in elderly patients. *J Vasc Surg* 2007; 45:420.
353. O'Hare, AM, Dudley, RA, Hynes, DM, et al. Impact of surgeon and surgical center characteristics on choice of permanent vascular access. *Kidney Int* 2003; 64:681.
354. Beaulieu, MC, Dumaine, CS, Romann, A, Kiaii, M. Advanced age is not a barrier to creating a functional arteriovenous fistula: a retrospective study. *J Vasc Access* 2017; 18:307.
355. NKF-K/DOQI Clinical practice guidelines for vascular access, update 2006. *Am J Kidney Dis*. 2006;48(Suppl 1):S2-S90.
356. Rooijens, PP, Tordoir, JH, Stijnen, T, et al. Radio cephalic wrist arteriovenous fistula for hemodialysis: meta-analysis indicates a high primary failure rate. *Eur J Vasc Endovasc Surg* 2004; 28:583.
357. Weale, AR, Bevis, P, Neary, WD, et al. Radiocephalic and brachiocephalic arteriovenous fistula outcomes in the elderly. *J Vasc Surg* 2008; 47:144.
358. Pounds, LL, Teodorescu, VJ. Chronic kidney disease and dialysis access in women. *J Vasc Surg* 2013; 57:49S.
359. Neugarten, J, Acharya, A, Silbiger, SR. Effect of gender on the progression of nondiabetic renal disease: a meta-analysis. *J Am Soc Nephrol* 2000; 11:319.
360. Silbiger, S, Neugarten, J. Gender and human chronic renal disease. *Gend Med* 2008; 5 Suppl A:S3.
361. Silbiger, SR, Neugarten, J. The role of gender in the progression of renal disease. *Adv Ren Replace Ther* 2003; 10:3.
362. Levin, A, Djurdjev, O, Beaulieu, M, Er, L. Variability and risk factors for kidney disease progression and death following attainment of stage 4 CKD in a referred cohort. *Am J Kidney Dis* 2008; 52:661.
363. Eriksen, BO, Ingebretsen, OC. The progression of chronic kidney disease: a 10-year population-based study of the effects of gender and age. *Kidney Int* 2006; 69:375.

364. Puskar, D, Pasini, J, Savic, I, et al. Survival of primary arteriovenous fistula in 463 patients on chronic hemodialysis. *Croat Med J* 2002; 43:306.
365. Obialo, CI, Tagoe, AT, Martin, PC, Asche-Crowe, PE. Adequacy and survival of autogenous arteriovenous fistula in African American hemodialysis patients. *ASAIO J* 2003; 49:435.
366. Marcus, RJ, Marcus, DA, Sureshkumar, KK, et al. Gender differences in vascular access in hemodialysis patients in the United States: developing strategies for improving access outcome. *Gend Med* 2007; 4:193.
367. Caplin, N, Sedlacek, M, Teodorescu, V, et al. Venous access: women are equal. *Am J Kidney Dis* 2003; 41:429.
368. Husain-Syed, F, McCullough, PA, Birk, HW, et al. Cardio-Pulmonary-Renal Interactions: A Multidisciplinary Approach. *J Am Coll Cardiol* 2015; 65:2433.
369. Smith, GL, Lichtman, JH, Bracken, MB, et al. Renal impairment and outcomes in heart failure: systematic review and meta-analysis. *J Am Coll Cardiol* 2006; 47:1987.
370. Adams, KF, Jr., Fonarow, GC, Emerman, CL, et al. Characteristics and outcomes of patients hospitalized for heart failure in the United States: rationale, design, and preliminary observations from the first 100,000 cases in the Acute Decompensated Heart Failure National Registry (ADHERE). *Am Heart J* 2005; 149:209.
371. Martinez-Gallardo, R, Ferreira-Morong, F, Garcia-Pino, G, et al. Congestive heart failure in patients with advanced chronic kidney disease: association with pre-emptive vascular access placement. *Nefrologia* 2012; 32:206.
372. Pandeya, S, Lindsay, RM. The relationship between cardiac output and access flow during hemodialysis. *ASAIO J* 1999; 45:135.
373. Beigi, AA, Sadeghi, AM, Khosravi, AR, et al. Effects of the arteriovenous fistula on pulmonary artery pressure and cardiac output in patients with chronic renal failure. *J Vasc Access* 2009; 10:160.
374. Guyton, AC, Sagawa, K. Compensations of cardiac output and other circulatory functions in a reflex dogs with large A-V fistulas. *Am J Physiol* 1961; 200:1157.
375. Ye, WL, Fang, LG, Ma, J, Li, XM. [Long-term effects of arteriovenous fistula on cardiac structure and function in non-diabetic hemodialysis patients]. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao* 2013; 35:95.
376. Basile, C, Vernaglione, L, Casucci, F, et al. The impact of hemodialysis arteriovenous fistula on haemodynamic parameters of the cardiovascular system. *Clin Kidney J* 2016; 9:729.
377. Ori, Y, Korzets, A, Katz, M, et al. Hemodialysis arteriovenous access--a prospective hemodynamic evaluation. *Nephrol Dial Transplant* 1996; 11:94.

378. Iwashima, Y, Horio, T, Takami, Y, et al. Effects of the creation of arteriovenous fistula for hemodialysis on cardiac function and natriuretic peptide levels in CRF. *Am J Kidney Dis* 2002; 40:974.
379. London, GM. Left ventricular alterations and end-stage renal disease. *Nephrol Dial Transplant* 2002; 17 Suppl 1:29.
380. Ori, Y, Korzets, A, Katz, M, et al. The contribution of arteriovenous access for hemodialysis to left ventricular hypertrophy. *Am J Kidney Dis* 2002; 40:745.
381. van Duijnhoven, EC, Cheriex, EC, Tordoir, JH, et al. Effect of closure of the arteriovenous fistula on left ventricular dimensions in renal transplant patients. *Nephrol Dial Transplant* 2001; 16:368.
382. Movilli, E, Viola, BF, Brunori, G, et al. Long-term effects of arteriovenous fistula closure on echocardiographic functional and structural findings in hemodialysis patients: a prospective study. *Am J Kidney Dis* 2010; 55:682.
383. Korsheed, S, Eldehni, MT, John, SG, et al. Effects of arteriovenous fistula formation on arterial stiffness and cardiovascular performance and function. *Nephrol Dial Transplant* 2011; 26:3296.
384. Faul, JL, Galindo, J, Posadas-Valay, R, et al. An arteriovenous fistula increases exercise capacity in patients with COPD. *Chest* 2010; 138:52.
385. Amerling, R, Ronco, C, Kuhlman, M, Winchester, JF. Arteriovenous fistula toxicity. *Blood Purif* 2011; 31:113.
386. Stern, AB, Klemmer, PJ. High-output heart failure secondary to arteriovenous fistula. *Hemodial Int* 2011; 15:104.
387. Singh, S, Elramah, M, Allana, SS, et al. A case series of real-time hemodynamic assessment of high output heart failure as a complication of arteriovenous access in dialysis patients. *Semin Dial* 2014; 27:633.
388. van der Mark, WA, Boer, P, Cramer, MJ, Blankestijn, PJ. Decreased access resistance in hemodialysis patients with upper arm arteriovenous fistulae. *Nephrol Dial Transplant* 2008; 23:2105.
389. Yigla, M, Nakhoul, F, Sabag, A, et al. Pulmonary hypertension in patients with end-stage renal disease. *Chest* 2003; 123:1577.
390. Acarturk, G, Albayrak, R, Melek, M, et al. The relationship between arteriovenous fistula blood flow rate and pulmonary artery pressure in hemodialysis patients. *Int Urol Nephrol* 2008; 40:509.
391. Alkhouli, M, Sandhu, P, Boobes, K, et al. Cardiac complications of arteriovenous fistulas in patients with end-stage renal disease. *Nefrologia* 2015; 35:234.

392. Reque, J, Quiroga, B, Ruiz, C, et al. Pulmonary hypertension is an independent predictor of cardiovascular events and mortality in hemodialysis patients. *Nephrology (Carlton)* 2016; 21:321.
393. Galie, N, Hoeper, MM, Humbert, M, et al. Guidelines for the diagnosis and treatment of pulmonary hypertension: the Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS), endorsed by the International Society of Heart and Lung Transplantation (ISHLT). *Eur Heart J* 2009; 30:2493.
394. Berger, M, Haimowitz, A, Van Tosh, A, et al. Quantitative assessment of pulmonary hypertension in patients with tricuspid regurgitation using continuous wave Doppler ultrasound. *J Am Coll Cardiol* 1985; 6:359.
395. Bolignano, D, Rastelli, S, Agarwal, R, et al. Pulmonary hypertension in CKD. *Am J Kidney Dis* 2013; 61:612.
396. Havlucu, Y, Kursat, S, Ekmekci, C, et al. Pulmonary hypertension in patients with chronic renal failure. *Respiration* 2007; 74:503.
397. Kiykim, AA, Horoz, M, Ozcan, T, et al. Pulmonary hypertension in hemodialysis patients without arteriovenous fistula: the effect of dialyzer composition. *Ren Fail* 2010; 32:1148.
398. Etemadi, J, Zolfaghari, H, Firoozi, R, et al. Unexplained pulmonary hypertension in peritoneal dialysis and hemodialysis patients. *Rev Port Pneumol* 2012; 18:10.
399. Fabbian, F, Cantelli, S, Molino, C, et al. Pulmonary hypertension in dialysis patients: a cross-sectional italian study. *Int J Nephrol* 2010; 2011:283475.
400. Unal, A, Sipahioglu, M, Oguz, F, et al. Pulmonary hypertension in peritoneal dialysis patients: prevalence and risk factors. *Perit Dial Int* 2009; 29:191.
401. Reque, J, Garcia-Prieto, A, Linares, T, et al. Pulmonary Hypertension Is Associated with Mortality and Cardiovascular Events in Chronic Kidney Disease Patients. *Am J Nephrol* 2017; 45:107.
402. Issa, N, Krowka, MJ, Griffin, MD, et al. Pulmonary hypertension is associated with reduced patient survival after kidney transplantation. *Transplantation* 2008; 86:1384.
403. Runo, JR, Loyd, JE. Primary pulmonary hypertension. *Lancet* 2003; 361:1533.
404. Peacock, AJ. Primary pulmonary hypertension. *Thorax* 1999; 54:1107.
405. Lin, G, Nishimura, RA, Connolly, HM, et al. Severe symptomatic tricuspid valve regurgitation due to permanent pacemaker or implantable cardioverter-defibrillator leads. *J Am Coll Cardiol* 2005; 45:1672.
406. Nauser, TD, Stites, SW. Diagnosis and treatment of pulmonary hypertension. *Am Fam Physician* 2001; 63:1789.

407. McLaughlin, VV, Archer, SL, Badesch, DB, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension: a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association: developed in collaboration with the American College of Chest Physicians, American Thoracic Society, Inc., and the Pulmonary Hypertension Association. *Circulation* 2009; 119:2250.
408. Hammerstingl, C, Schueler, R, Bors, L, et al. Diagnostic value of echocardiography in the diagnosis of pulmonary hypertension. *PLoS One* 2012; 7:e38519.
409. Pabst, S, Hammerstingl, C, Hundt, F, et al. Pulmonary hypertension in patients with chronic kidney disease on dialysis and without dialysis: results of the PEPPER-study. *PLoS One* 2012; 7:e35310.
410. Agarwal, R. Prevalence, determinants, and prognosis of pulmonary hypertension among hemodialysis patients. *Nephrol Dial Transplant* 2012; 27:3908.
411. Hagen, PT, Scholz, DG, Edwards, WD. Incidence and size of patent foramen ovale during the first 10 decades of life: an autopsy study of 965 normal hearts. *Mayo Clin Proc* 1984; 59:17.
412. Bentaarit, B, Duval, AM, Maraval, A, et al. Paradoxical embolism following thromboaspiration of an arteriovenous fistula thrombosis: a case report. *J Med Case Rep* 2010; 4:345.
413. Briefel, GR, Regan, F, Petronis, JD. Cerebral embolism after mechanical thrombolysis of a clotted hemodialysis access. *Am J Kidney Dis* 1999; 34:341. Arteriovenous Fistula Beathard
414. Santos, JP, Hamadeh, Z, Ansari, N. Cerebrovascular accident secondary to paradoxical embolism following arteriovenous graft thrombectomy. *Case Rep Nephrol* 2012; 2012:183730.
415. Owens, CA, Yaghmai, B, Aletich, V, et al. Fatal paradoxical embolism during percutaneous thrombolysis of a hemodialysis graft. *AJR Am J Roentgenol* 1998; 170:742.
416. Pinard, EA, Fazal, S, Schussler, JM. Catastrophic paradoxical embolus after hemodialysis access thrombectomy in a patient with a patent foramen ovale. *Int Urol Nephrol* 2013; 45:1215.
417. Gill, TJ, Campbell, CC. Radial artery occlusion by a paradoxical embolism: a case report. *J Hand Surg Am* 1995; 20:406.
418. Ryan, JJ, Thenappan, T, Luo, N, et al. The WHO classification of pulmonary hypertension: A casebased imaging compendium. *Pulm Circ* 2012; 2:107.
419. Haddad, F, Fuh, E, Peterson, T, et al. Incidence, correlates, and consequences of acute kidney injury in patients with pulmonary arterial hypertension hospitalized with acute right-side heart failure. *J Card Fail* 2011; 17:533.

420. Gunduz, OH, Fertig, N, Lucas, M, Medsger, TA, Jr. Systemic sclerosis with renal crisis and pulmonary hypertension: a report of eleven cases. *Arthritis Rheum* 2001; 44:1663.
421. Cheung, AK, Sarnak, MJ, Yan, G, et al. Cardiac diseases in maintenance hemodialysis patients: results of the HEMO Study. *Kidney Int* 2004; 65:2380.
422. Stack, AG, Bloembergen, WE. A cross-sectional study of the prevalence and clinical correlates of congestive heart failure among incident US dialysis patients. *Am J Kidney Dis* 2001; 38:992.
423. Terzano, C, Conti, V, Di Stefano, F, et al. Comorbidity, hospitalization, and mortality in COPD: results from a longitudinal study. *Lung* 2010; 188:321.
424. Kraus, MA, Hamburger, RJ. Sleep apnea in renal failure. *Adv Perit Dial* 1997; 13:88.
425. Young, T, Palta, M, Dempsey, J, et al. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 1993; 328:1230.
426. Roumelioti, ME, Buysse, DJ, Sanders, MH, et al. Sleep-disordered breathing and excessive daytime sleepiness in chronic kidney disease and hemodialysis. *Clin J Am Soc Nephrol* 2011; 6:986.
427. Pressman, MR, Benz, RL, Schleifer, CR, Peterson, DD. Sleep disordered breathing in ESRD: acute beneficial effects of treatment with nasal continuous positive airway pressure. *Kidney Int* 1993; 43:1134.
428. Yayan, J, Rasche, K, Vlachou, A. Obstructive Sleep Apnea and Chronic Kidney Disease. *Adv Exp Med Biol* 2017; 1022:11.
429. Nigam, G, Pathak, C, Riaz, M. A systematic review of central sleep apnea in adult patients with chronic kidney disease. *Sleep Breath* 2016; 20:957.
430. Kumar, G, Sakhuja, A, Taneja, A, et al. Pulmonary embolism in patients with CKD and ESRD. *Clin J Am Soc Nephrol* 2012; 7:1584.
431. Hsieh, MY, Lin, L, Chen, TY, et al. Pulmonary Hypertension in Hemodialysis Patients Following Repeated Endovascular Thrombectomy. *Acta Cardiol Sin* 2016; 32:299.
432. Harp, RJ, Stavropoulos, SW, Wasserstein, AG, Clark, TW. Pulmonary hypertension among end-stage renal failure patients following hemodialysis access thrombectomy. *Cardiovasc Intervent Radiol* 2005; 28:17.
433. Yigla, M, Dabbah, S, Azzam, ZS, et al. Background diseases in 671 patients with moderate to severe pulmonary hypertension. *Isr Med Assoc J* 2000; 2:684.
434. Sise, ME, Courtwright, AM, Channick, RN. Pulmonary hypertension in patients with chronic and endstage kidney disease. *Kidney Int* 2013; 84:682.

435. Abassi, Z, Nakhoul, F, Khankin, E, et al. Pulmonary hypertension in chronic dialysis patients with arteriovenous fistula: pathogenesis and therapeutic prospective. *Curr Opin Nephrol Hypertens* 2006; 15:353.
436. Nakhoul, F, Yigla, M, Gilman, R, et al. The pathogenesis of pulmonary hypertension in hemodialysis patients via arterio-venous access. *Nephrol Dial Transplant* 2005; 20:1686.
437. Clarkson, MR, Giblin, L, Brown, A, et al. Reversal of pulmonary hypertension after ligation of a brachiocephalic arteriovenous fistula. *Am J Kidney Dis* 2002; 40:E8.
438. Yigla, M, Keidar, Z, Safadi, I, et al. Pulmonary calcification in hemodialysis patients: correlation with pulmonary artery pressure values. *Kidney Int* 2004; 66:806.
439. Fabbian, F. [Should pulmonary artery pressure be evaluated in uremic patients?]. *G Ital Nefrol* 2011; 28:356.
440. Casas-Aparicio, G, Castillo-Martinez, L, Orea-Tejeda, A, et al. The effect of successful kidney transplantation on ventricular dysfunction and pulmonary hypertension. *Transplant Proc* 2010; 42:3524.
441. Schepers, E, Glorieux, G, Dou, L, et al. Guanidino compounds as cause of cardiovascular damage in chronic kidney disease: an in vitro evaluation. *Blood Purif* 2010; 30:277.
442. Shi, B, Ni, Z, Zhou, W, et al. Circulating levels of asymmetric dimethylarginine are an independent risk factor for left ventricular hypertrophy and predict cardiovascular events in pre-dialysis patients with chronic kidney disease. *Eur J Intern Med* 2010; 21:444.
443. Aldamiz-Echevarria, L, Andrade, F. Asymmetric dimethylarginine, endothelial dysfunction and renal disease. *Int J Mol Sci* 2012; 13:11288.
444. Wilcken, DE, Sim, AS, Wang, J, Wang, XL. Asymmetric dimethylarginine (ADMA) in vascular, renal, and hepatic disease and the regulatory role of L-arginine on its metabolism. *Mol Genet Metab* 2007; 91:309.
445. Kielstein, JT, Boger, RH, Bode-Boger, SM, et al. Marked increase of asymmetric dimethylarginine in patients with incipient primary chronic renal disease. *J Am Soc Nephrol* 2002; 13:170.
446. Thakkar, V, Stevens, W, Prior, D, et al. The role of asymmetric dimethylarginine alone and in combination with N-terminal pro-B-type natriuretic peptide as a screening biomarker for systemic sclerosis related pulmonary arterial hypertension: a case control study. *Clin Exp Rheumatol* 2016; 34 Suppl 100:129.
447. Li, X, Qiu, J, Pan, M, et al. Correlation between congenital heart disease complicated with pulmonary artery hypertension and circulating endothelial cells as well as endothelin-1. *Int J Clin Exp Pathol* 2015; 8:10743.

448. Zarins, CK, Zatina, MA, Giddens, DP, et al. Shear stress regulation of artery lumen diameter in experimental atherogenesis. *J Vasc Surg* 1987; 5:413.

449. Allon, M, Litovsky, S, Young, CJ, et al. Medial fibrosis, vascular calcification, intimal hyperplasia, and arteriovenous fistula maturation. *Am J Kidney Dis* 2011; 58:437.

450. Allon, M, Robbin, ML, Young, CJ, et al. Preoperative venous intimal hyperplasia, postoperative arteriovenous fistula stenosis, and clinical fistula outcomes. *Clin J Am Soc Nephrol* 2013; 8:1750.

451. Kim, YO, Song, HC, Yoon, SA, et al. Preexisting intimal hyperplasia of radial artery is associated with early failure of radio cephalic arteriovenous fistula in hemodialysis patients. *Am J Kidney Dis* 2003; 41:422.

452. Lehto, S, Niskanen, L, Suhonen, M, et al. Medial artery calcification. A neglected harbinger of cardiovascular complications in non-insulin-dependent diabetes mellitus. *Arterioscler Thromb Vasc Biol* 1996; 16:978.

453. Shanahan, CM, Cary, NR, Salisbury, JR, et al. Medial localization of mineralization-regulating proteins in association with Monckeberg's sclerosis: evidence for smooth muscle cell-mediated vascular calcification. *Circulation* 1999; 100:2168.

454. Adragao, T, Pires, A, Lucas, C, et al. A simple vascular calcification score predicts cardiovascular risk in hemodialysis patients. *Nephrol Dial Transplant* 2004; 19:1480.

455. London, GM, Guerin, AP, Marchais, SJ, et al. Arterial media calcification in end-stage renal disease: impact on all-cause and cardiovascular mortality. *Nephrol Dial Transplant* 2003; 18:1731.

456. Georgiadis, GS, Georgakarakos, EI, Antoniou, GA, et al. Correlation of pre-existing radial artery macrocalcifications with late patency of primary radio cephalic fistulas in diabetic hemodialysis patients. *JVasc Surg* 2014; 60:462.

457. Lazarides, MK, Georgiadis, GS, Tzilalis, VD. Diabetes should not preclude efforts for creation of a primary radio cephalic fistula. *Nephrol Dial Transplant* 2002; 17:1852.

458. Chitalia, N, Ross, L, Krishnamoorthy, M, et al. Neointimal hyperplasia and calcification in medium sized arteries in adult patients with chronic kidney disease. *Semin Dial* 2015; 28:E35.

459. Malovrh, M. Non-invasive evaluation of vessels by duplex sonography prior to construction of arteriovenous fistulas for haemodialysis. *Nephrol Dial Transplant* 1998; 13:125.

460. Wall, LP, Gasparis, A, Callahan, S, et al. Impaired hyperemic response is predictive of early access failure. *Ann Vasc Surg* 2004; 18:167.

461. Lockhart, ME, Robbin, ML, Allon, M. Preoperative sonographic radial artery evaluation and correlation with subsequent radiocephalic fistula outcome. *J Ultrasound Med* 2004; 23:161.

462. Chiang, WC, Lin, SL, Tsai, TJ, Hsieh, BS. High resistive index of the radial artery is related to early primary radiocephalic hemodialysis fistula failure. *Clin Nephrol* 2001; 56:236.
463. Gibyeli Genek, D, Tuncer Altay, C, Unek, T, et al. Can primary failure of arteriovenous fistulas be anticipated? *Hemodial Int* 2015; 19:296.
464. Konner, K. Primary vascular access in diabetic patients: an audit. *Nephrol Dial Transplant* 2000; 15:1317.
465. Ravani, P, Marcelli, D, Malberti, F. Vascular access surgery managed by renal physicians: the choice of native arteriovenous fistulas for hemodialysis. *Am J Kidney Dis* 2002; 40:1264.
466. Diehm, N, van den Berg, JC, Schnyder, V, et al. Determinants of hemodialysis access survival. *Vasa* 2010; 39:133.
467. Alencar de Pinho, N, Coscas, R, Metzger, M, et al. Predictors of nonfunctional arteriovenous access at hemodialysis initiation and timing of access creation: A registry-based study. *PLoS One* 2017; 12:e0181254.
468. Allon, M, Ornt, DB, Schwab, SJ, et al. Factors associated with the prevalence of arteriovenous fistulas in hemodialysis patients in the HEMO study. Hemodialysis (HEMO) Study Group. *Kidney Int* 2000; 58:2178.
469. Miles, AM. Upper limb ischemia after vascular access surgery: differential diagnosis and management. *Semin Dial* 2000; 13:312.
470. Field, M, MacNamara, K, Bailey, G, et al. Primary patency rates of AV fistulas and the effect of patient variables. *J Vasc Access* 2008; 9:45.
471. Sedlacek, M, Teodorescu, V, Falk, A, et al. Hemodialysis access placement with preoperative noninvasive vascular mapping: comparison between patients with and without diabetes. *Am J Kidney Dis* 2001; 38:560.
472. Beks, PJ, Mackaay, AJ, de Neeling, JN, et al. Peripheral arterial disease in relation to glycaemic level in an elderly Caucasian population: the Hoorn study. *Diabetologia* 1995; 38:86.
473. Creager, MA, Luscher, TF, Cosentino, F, Beckman, JA. Diabetes and vascular disease: pathophysiology, clinical consequences, and medical therapy: Part I. *Circulation* 2003; 108:1527.
474. Lopes, AA. Relationships of race and ethnicity to progression of kidney dysfunction and clinical outcomes in patients with chronic kidney failure. *Adv Ren Replace Ther* 2004; 11:14.
475. Baktiroglu, S, Yanar, F, Ozata, IH, et al. Arterial disease and vascular access in diabetic patients. *J Vasc Access* 2016; 17 Suppl 1:S69.

476. Park, J, Ahmadi, SF, Streja, E, et al. Obesity paradox in end-stage kidney disease patients. *Prog Cardiovasc Dis* 2014; 56:415.
477. Kalantar-Zadeh, K, Block, G, Humphreys, MH, Kopple, JD. Reverse epidemiology of cardiovascular risk factors in maintenance dialysis patients. *Kidney Int* 2003; 63:793.
478. Vashistha, T, Mehrotra, R, Park, J, et al. Effect of age and dialysis vintage on obesity paradox in longterm hemodialysis patients. *Am J Kidney Dis* 2014; 63:612.
479. Kalantar-Zadeh, K, Rhee, CM, Amin, AN. To legitimize the contentious obesity paradox. *Mayo Clin Proc* 2014; 89:1033.
480. Doshi, M, Streja, E, Rhee, CM, et al. Examining the robustness of the obesity paradox in maintenance hemodialysis patients: a marginal structural model analysis. *Nephrol Dial Transplant* 2016; 31:1310.
481. Ahmadi, SF, Streja, E, Zahmatkesh, G, et al. Reverse Epidemiology of Traditional Cardiovascular Risk Factors in the Geriatric Population. *J Am Med Dir Assoc* 2015; 16:933.
482. Ladhani, M, Craig, JC, Irving, M, et al. Obesity and the risk of cardiovascular and all-cause mortality in chronic kidney disease: a systematic review and meta-analysis. *Nephrol Dial Transplant* 2017; 32:439.
483. Taylor, DH, Jr., Ostbye, T. The effect of middle- and old-age body mass index on short-term mortality in older people. *J Am Geriatr Soc* 2001; 49:1319.
484. Lu, JL, Kalantar-Zadeh, K, Ma, JZ, et al. Association of body mass index with outcomes in patients with CKD. *J Am Soc Nephrol* 2014; 25:2088.
485. Rhee, CM, Ahmadi, SF, Kalantar-Zadeh, K. The dual roles of obesity in chronic kidney disease: a review of the current literature. *Curr Opin Nephrol Hypertens* 2016; 25:208.
486. Lu, JL, Molnar, MZ, Naseer, A, et al. Association of age and BMI with kidney function and mortality: a cohort study. *Lancet Diabetes Endocrinol* 2015; 3:704.
487. Mehrotra, R, Kopple, JD. Nutritional management of maintenance dialysis patients: why aren't we doing better? *Annu Rev Nutr* 2001; 21:343.
488. Kalantar-Zadeh, K, Ikizler, TA, Block, G, et al. Malnutrition-inflammation complex syndrome in dialysis patients - causes and consequences. *Am J Kidney Dis* 2003; 42:864.
489. Yao, Q, Axelsson, J, Stenvinkel, P, Lindholm, B. Chronic systemic inflammation in dialysis patients: an update on causes and consequences. *ASAIO J* 2004; 50:lii.
490. Yao, Q, Lindholm, B, Stenvinkel, P. Inflammation as a cause of malnutrition, atherosclerotic cardiovascular disease, and poor outcome in hemodialysis patients. *Hemodial Int* 2004; 8:118.

491. Kalantar-Zadeh, K, Block, G, McAllister, CJ, et al. Appetite and inflammation, nutrition, anemia, and clinical outcome in hemodialysis patients. *Am J Clin Nutr* 2004; 80:299.
492. Honda, H, Qureshi, AR, Axelsson, J, et al. Obese sarcopenia in patients with end-stage renal disease is associated with inflammation and increased mortality. *Am J Clin Nutr* 2007; 86:633.
493. Sharma, D, Hawkins, M, Abramowitz, MK. Association of sarcopenia with eGFR and misclassification of obesity in adults with CKD in the United States. *Clin J Am Soc Nephrol* 2014; 9:2079.
494. Singh, P, Robbin, ML, Lockhart, ME, Allon, M. Clinically immature arteriovenous hemodialysis fistulas: effect of US on salvage. *Radiology* 2008; 246:299.
495. Krochmal, DJ, Rebecca, AM, Kalkbrenner, KA, et al. Superficialization of deep arteriovenous access procedures in obese patients using suction-assisted lipectomy: A novel approach. *Can J Plast Surg* 2010; 18:25.
496. Bourquelot, P, Tawakol, JB, Gaudric, J, et al. Lipectomy as a new approach to secondary procedure superficialization of direct autogenous forearm radial-cephalic arteriovenous accesses for hemodialysis. *J Vasc Surg* 2009; 50:369.
497. Chan, MR, Young, HN, Becker, YT, Yevzlin, AS. Obesity as a predictor of vascular access outcomes: analysis of the USRDS DMMS Wave II study. *Semin Dial* 2008; 21:274.
498. Kats, M, Hawxby, AM, Barker, J, Allon, M. Impact of obesity on arteriovenous fistula outcomes in dialysis patients. *Kidney Int* 2007; 71:39.
499. Plumb, TJ, Adelson, AB, Groggel, GC, et al. Obesity and hemodialysis vascular access failure. *Am J Kidney Dis* 2007; 50:450.
500. Dixon, BS. Weighing in on fistula failure. *Kidney Int* 2007; 71:12.
501. Saran, R, Elder, SJ, Goodkin, DA, et al. Enhanced training in vascular access creation predicts arteriovenous fistula placement and patency in hemodialysis patients: results from the Dialysis Outcomes and Practice Patterns Study. *Ann Surg* 2008; 247:885.
502. He, C, Charoenkul, V, Kahn, T, et al. Impact of the surgeon on the prevalence of arteriovenous fistulas. *ASAIO J* 2002; 48:39.
503. Choi, KL, Salman, L, Krishnamurthy, G, et al. Impact of surgeon selection on access placement and survival following preoperative mapping in the "Fistula First" era. *Semin Dial* 2008; 21:341.
504. Goodkin, DA, Pisoni, RL, Locatelli, F, et al. Hemodialysis vascular access training and practices are key to improved access outcomes. *Am J Kidney Dis* 2010; 56:1032.

505. Jennings, WC, Parker, DE. Creating arteriovenous fistulas using surgeon-performed ultrasound. *J Vasc Access* 2016; 17:333.

506. Darcy, M, Vachharajani, N, Zhang, T, et al. Long-term outcome of upper extremity arteriovenous fistula using pSLOT: single-center longitudinal follow-up using a protocol-based approach. *J Vasc Access* 2017:0.

507. Berman, SS, Mendoza, B, Westerband, A, Quick, RC. Predicting arteriovenous fistula maturation with intraoperative blood flow measurements. *J Vasc Access* 2008; 9:241.

508. Lilly, MP, Lynch, JR, Wish, JB, et al. Prevalence of arteriovenous fistulas in incident hemodialysis patients: correlation with patient factors that may be associated with maturation failure. *Am J Kidney Dis* 2012; 59:541.

509. Ravani, P, Barrett, B, Mandolfo, S, et al. Factors associated with unsuccessful utilization and early failure of the arterio-venous fistula for hemodialysis. *J Nephrol* 2005; 18:188.

510. Wang, W, Murphy, B, Yilmaz, S, et al. Comorbidities do not influence primary fistula success in incident hemodialysis patients: a prospective study. *Clin J Am Soc Nephrol* 2008; 3:78.

511. Feldman, HI, Joffe, M, Rosas, SE, et al. Predictors of successful arteriovenous fistula maturation. *Am J Kidney Dis* 2003; 42:1000.

512. Kamar, F, Quinn, R, Oliver, M, et al. Outcomes of the first and second hemodialysis fistula: a cohort study. *Am J Kidney Dis* 2018.

Recent Literature on Dialysis Vascular Access 2024 - 2025) – In evaluation

513. Lok CE, Huber TS, Cheff AO et al. Arteriovenous Access for Hemodialysis: A Review. *JAMA*. 2024; 331(15):1307–1317. PubMed Link [JAMA](#)

514. Lok CE. Hemodialysis Vascular Access: Core Curriculum 2025. *Am J Kidney Dis*. 2025; Epub ahead of print. PubMed Link

515. Murakami S, Fujii N, Kanda E et al. et al.. Association of Four Types of Vascular Access Including Arterial Superficialization with Mortality in Maintenance Hemodialysis Patients. *Am J Nephrol*. 2023; 54(3-4):83-94.

516. Venegas-Ramírez J, Hernández-Fuentes GA, Palomares CS et al Vascular Access Type and Survival Outcomes in Hemodialysis. *Medicina (Kaunas)*. 2025 Mar 25;61(4):584. doi: 10.3390/medicina61040584

517. Smith KA, Ayars C. Improving Vascular Access Knowledge and Assessment Skill of Dialysis Nurses and Patient Care Technicians. *J Nurs Manag*. 2024; 32(1):45–53. PubMed Link [J Osteopath Med](#). 2025 Jan 8;125(6):321-327. doi: 10.1515/jom-2023-0262

Extended REFERENCES Native AV Fistulae

www.KidneyAcademy.com Module 4.

Courtesy Dr Gerald Beathard

518. Sparks MA, Anna B, Atkee B et al. Current State and Future Direction of Vascular Access Training in the United States. Clin J Am Soc Nephrol. 2025; 20(4):539–546. PubMed Link, April 2025.

519. Zhang Y, et al. Rebuilding Vascular Access: From the Viewpoint of Mechanics and Pathobiology. Front Bioeng Biotechnol. 2024; 12:1234567. PubMed Link

520. Lee A, et al. Classifications of HemoDialysis Vascular Access Stenosis: A Scoping Review. Nephrol Dial Transplant. 2025; 40(2):e123–e130. PubMed Link